

The Two Rigid Body Interaction using Angular Momentum Theory Formulae

Gwenaël Boué

► **To cite this version:**

Gwenaël Boué. The Two Rigid Body Interaction using Angular Momentum Theory Formulae. *Celestial Mechanics and Dynamical Astronomy*, Springer Verlag, 2016, 128 (2-3), pp.261-273. <10.1007/s10569-017-9751-2>. <hal-01465812>

HAL Id: hal-01465812

<http://hal.upmc.fr/hal-01465812>

Submitted on 14 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Two Rigid Body Interaction using Angular Momentum Theory Formulae

Gwenaél Boué

Received: date / Accepted: date

Abstract This work presents an elegant formalism to model the evolution of the full two rigid body problem. The equations of motion, given in a Cartesian coordinate system, are expressed in terms of spherical harmonics and Wigner D-matrices. The algorithm benefits from the numerous recurrence relations satisfied by these functions allowing a fast evaluation of the mutual potential. Moreover, forces and torques are straightforwardly obtained by application of ladder operators taken from the angular momentum theory and commonly used in quantum mechanics. A numerical implementation of this algorithm is made. Tests show that the present code is significantly faster than those currently available in literature.

Keywords full two rigid problem · binary systems · spin-orbit coupling · numerical method

1 Introduction

Modelling the evolution of two rigid bodies with arbitrary shapes in gravitational interaction is not an easy task. The first obstacle is the determination of the mutual potential, the second is the derivation of the equations of motion in a suitable form to allow fast computation. In general cases, the potential has to be expanded and truncated at some order in the ratio of the bodies' mean radii to the distance between the two barycenters. Using the angular momentum theory developed by [Wigner \(1959\)](#), [Borderies \(1978\)](#) manage to provide a compact expression of the mutual potential at any order. In this expression, the gravity field of each body is described by Stokes coefficients, the relative orientation of the two bodies appears through Wigner D-matrices – also called Euler functions ([Borderies, 1978](#)) – and the dependence in the distance between the two barycenters is embedded in solid harmonics. Equations of motion associated to this expansion have been proposed by [Maciejewski \(1995\)](#). A strictly equivalent formalism based on symmetric trace-free (STF) tensor ([Hartmann et al, 1994](#); [Mathis and Le Poncin-Lafitte, 2009](#)) has

G. Boué
E-mail: gwenael.boue@obspm.fr IMCCE, Observatoire de Paris, UPMC Univ. Paris 6, PSL Research University, Paris, France

been implemented by [Compère and Lemaître \(2014\)](#) who studied numerically the evolution of the binary asteroid 1999 KW4.

The decomposition in spherical harmonics has sometimes been discarded based on the misconception that this formalism involves many trigonometric functions increasing as much the computation time and the risk of numerical instabilities. To circumvent this issue, alternative decompositions of the potential have been applied. For instance, [Paul \(1988\)](#) explicitly wrote the mutual potential in Cartesian coordinates at all orders. The associated equations of motion were then provided by [Tricarico \(2008\)](#). Recently, [Hou et al \(2016\)](#) revisited this approach and built the as yet fastest algorithm able to integrate the full two rigid body problem thanks to a set of recurrence formulae whose coefficients can be computed and stored beforehand. Another example of alternative is the so-called polyhedron approach relying on Stokes' theorem in which the volume integral leading to the mutual potential is converted into a surface integral over the boundaries of the two bodies ([Werner and Scheeres, 2005](#)). This algorithm implemented by [Fahnstock and Scheeres \(2006\)](#) was at the time the fastest integrator.

Here we revisit [Borderies'](#) expression of the mutual potential developed in spherical harmonics, but we express them in Cartesian coordinates. By consequence, the description of the orbital configuration is actually equivalent to those where the potential is explicitly written in Cartesian coordinates as in ([Paul, 1988](#)), for instance. The main contribution of the present study is in the parametrisation of rotations. Whereas in previous works, torques are computed by an explicit derivation of the potential energy with respect to Euler angles or with respect to rotation matrix elements, here forces and torques are simply obtained by application of ladder operators taken from quantum mechanics theory. This approach has already been successfully applied to the modelling of tidal evolution of close-in planets ([Boué et al, 2016](#)). In this study we show that it allows to build the fastest numerical integrator of the full two rigid body problem.

The paper is organised as follows: in Section 2, the equations of motion are computed using Poincaré's method ([Poincaré, 1901](#)). Forces and torques are written in terms of ladder operators commonly used in quantum mechanics. The section also contains all the recurrence relations allowing an efficient numerical implementation of the problem. Numerical tests are performed in Section 3. Conclusions are drawn in the last section.

2 Equations of motion

Let two rigid bodies A and B with arbitrary shapes in gravitational interaction in an inertial frame $(O; \mathbf{e}_x, \mathbf{e}_y, \mathbf{e}_z)$. For both bodies A and B we define body-fixed frames centred on their respective barycenters O_A and O_B which are $(O_A; \mathbf{e}_x^A, \mathbf{e}_y^A, \mathbf{e}_z^A)$ and $(O_B; \mathbf{e}_x^B, \mathbf{e}_y^B, \mathbf{e}_z^B)$, respectively. To fasten the integration, the problem is not described in the inertial frame as in ([Borderies, 1978](#)), but in the body-fixed frame of A ([Maciejewski, 1995](#)). This is also the choice of [Hou et al \(2016\)](#) with whom we wish to compare the method. The convention used in this paper is the following: unprimed quantities are expressed in the body-fixed frame of A , while primed ones are written in the body-fixed frame of B . Vectors expressed in the inertial frame are written with the superscript 0.

2.1 Lagrangian

Following [Maciejewski \(1995\)](#), the potential reads

$$U_{AB} = -GM_A M_B \sum_{l_1=0}^{\infty} \sum_{m_1=-l_1}^{l_1} \sum_{l_2=0}^{\infty} \sum_{m_2=-l_2}^{l_2} \left(R_A^{l_1} R_B^{l_2} Z_{l_1, m_1}^A Z_{l_2, m_2}^B \right. \\ \left. \times (-1)^{l_2} \gamma_{l_2, m_2}^{l_1, m_1} \frac{Y_{l_1+l_2, m_1+m_2}(\mathbf{r})}{r^{l_1+l_2+1}} \right) \quad (1)$$

where M_* , R_* , and $Z_{l,m}^*$ are the mass, the mean radius and Stokes coefficients of the body $* = A, B$, respectively. Note that $Z_{l,m}^A$ is constant while $Z_{l,m}^B$ is not because both quantities are written in the body-fixed frame of A . The radius vector $\mathbf{r} = \mathbf{r}_B - \mathbf{r}_A$ connects the two barycenters. $Y_{l,m}$ are complex spherical harmonics and $\gamma_{l_2, m_2}^{l_1, m_1}$ are constant coefficients. Here we use the Schmidt semi-normalisation of the spherical harmonics, such that,

$$Y_{l,m}(\theta, \phi) = (-1)^m \sqrt{\frac{(l-m)!}{(l+m)!}} P_{l,m}(\cos \theta) e^{im\phi} \quad (2)$$

where the associated Legendre polynomials $P_{l,m}$ are defined as

$$P_{l,m}(x) = \frac{1}{2^l l!} (1-x^2)^{m/2} \frac{d^{l+m}}{dx^{l+m}} (x^2-1)^l.$$

The complex Stokes coefficients $Z_{l,m}$ are related to the usual real Stokes coefficients $C_{l,m}$ and $S_{l,m}$ by

$$Z_{l,m} = (-1)^m \frac{1 + \delta_{m,0}}{2} \sqrt{\frac{(l+m)!}{(l-m)!}} (C_{l,m} - iS_{l,m}), \quad \text{for } m \geq 0 \quad (3a)$$

and the symmetry relation

$$Z_{l,m} = (-1)^m \bar{Z}_{l,-m}, \quad \text{for } m < 0. \quad (3b)$$

In Eq. (3a), $\delta_{i,j}$ is the Kronecker delta equal to 1 if $i = j$ and equal to 0 otherwise. In Eq. (3b), the bar above $Z_{l,m}$ denotes the complex conjugate. Following our normalisation convention, the coefficients $\gamma_{l_2, m_2}^{l_1, m_1}$ are

$$\gamma_{l_2, m_2}^{l_1, m_1} = \sqrt{\frac{(l_1 + l_2 - m_1 - m_2)! (l_1 + l_2 + m_1 + m_2)!}{(l_1 + m_1)! (l_1 - m_1)! (l_2 + m_2)! (l_2 - m_2)!}}.$$

Let

$$\mathbf{C}_A = [\mathbf{e}_x^A, \mathbf{e}_y^A, \mathbf{e}_z^A], \quad \mathbf{C}_B = [\mathbf{e}_x^B, \mathbf{e}_y^B, \mathbf{e}_z^B] \quad \text{and} \quad \mathbf{C} = \mathbf{C}_A^T \mathbf{C}_B$$

be the rotation matrices such that

$$\mathbf{r} = \mathbf{C}_A^T \mathbf{r}^0, \quad Z_{l,m}^B = \sum_{m'=-l}^l D_{m,m'}^l(\mathbf{C}) Z_{l,m'}^B,$$

where \mathbf{C}_A^T means the transpose of \mathbf{C}_A and $D_{m,m'}^l(\mathbf{C})$ is an element of Wigner D-matrix associated to the rotation \mathbf{C} . $Z_{l,m'}^B$ are the (constant) complex Stokes coefficients of body B expressed in its body-fixed frame.

In the barycentric frame, the kinetic energy of the system reads

$$T = \frac{1}{2} \boldsymbol{\Omega}_A^T \mathbf{I}_A \boldsymbol{\Omega}_A + \frac{1}{2} (\boldsymbol{\Omega} + \boldsymbol{\Omega}_A)^T \mathbf{I}_B (\boldsymbol{\Omega} + \boldsymbol{\Omega}_A) + \frac{1}{2} \mu \|\mathbf{v} + \boldsymbol{\Omega}_A \times \mathbf{r}\|^2 \quad (4)$$

where \mathbf{I}_A and \mathbf{I}_B are the inertia matrices of the bodies A and B expressed in the body-fixed frame of A , respectively. $\boldsymbol{\Omega}_A$ is the rotation vector of the body A with respect to the inertial frame, while $\boldsymbol{\Omega} = \boldsymbol{\Omega}_B - \boldsymbol{\Omega}_A$ is the rotation vector of the body B relative to the body A . $\mu = M_A M_B / (M_A + M_B)$ is the reduced mass and $\mathbf{v} = \dot{\mathbf{r}}$ is the relative velocity in the body-fixed frame of A . In the expression of the kinetic energy (Eq. 4), \mathbf{I}_A is constant while $\mathbf{I}_B = \mathbf{C} \mathbf{I}_B' \mathbf{C}^T$ is a function of \mathbf{C} .

The Lagrangian of the problem $f = T - U_{AB}$ is a function of $\mathbf{q} = (\mathbf{C}_A, \mathbf{C}, \mathbf{r})$ and $\boldsymbol{\eta} = (\boldsymbol{\Omega}_A, \boldsymbol{\Omega}, \mathbf{v})$. Note that \mathbf{C}_A and \mathbf{C} are themselves functions of only three coordinates, such as the 3-2-3 Euler angles $(\alpha_A, \beta_A, \gamma_A)$ and (α, β, γ) , respectively:

$$\mathbf{C}_A(\alpha_A, \beta_A, \gamma_A) = \mathbf{C}_z(\alpha_A) \mathbf{C}_y(\beta_A) \mathbf{C}_z(\gamma_A), \quad \mathbf{C}(\alpha, \beta, \gamma) = \mathbf{C}_z(\alpha) \mathbf{C}_y(\beta) \mathbf{C}_z(\gamma).$$

To retrieve the equations of motion given by [Maciejewski \(1995\)](#), we use Poincaré's *forme nouvelle des équations de la mécanique* ([Poincaré, 1901](#)). We define a vector field basis associated to infinitesimal rotations of the body A as

$$\hat{\mathcal{J}}_{A,x} = \cos \gamma_A \cot \beta_A \frac{\partial}{\partial \gamma_A} + \sin \gamma_A \frac{\partial}{\partial \beta_A} - \frac{\cos \gamma_A}{\sin \beta_A} \frac{\partial}{\partial \alpha_A}, \quad (5a)$$

$$\hat{\mathcal{J}}_{A,y} = -\sin \gamma_A \cot \beta_A \frac{\partial}{\partial \gamma_A} + \cos \gamma_A \frac{\partial}{\partial \beta_A} + \frac{\sin \gamma_A}{\sin \beta_A} \frac{\partial}{\partial \alpha_A}, \quad (5b)$$

$$\hat{\mathcal{J}}_{A,z} = \frac{\partial}{\partial \gamma_A}. \quad (5c)$$

Equivalently, we introduce a basis field corresponding to infinitesimal rotations of the body B relative to the body A :

$$\hat{\mathcal{J}}_x = -\cos \alpha \cot \beta \frac{\partial}{\partial \alpha} - \sin \alpha \frac{\partial}{\partial \beta} + \frac{\cos \alpha}{\sin \beta} \frac{\partial}{\partial \gamma}, \quad (6a)$$

$$\hat{\mathcal{J}}_y = -\sin \alpha \cot \beta \frac{\partial}{\partial \alpha} + \cos \alpha \frac{\partial}{\partial \beta} + \frac{\sin \alpha}{\sin \beta} \frac{\partial}{\partial \gamma}, \quad (6b)$$

$$\hat{\mathcal{J}}_z = \frac{\partial}{\partial \alpha}. \quad (6c)$$

At last, we consider the canonical vector field basis associated to infinitesimal translations

$$\hat{\mathcal{P}}_x = \frac{\partial}{\partial x}, \quad (7a)$$

$$\hat{\mathcal{P}}_y = \frac{\partial}{\partial y}, \quad (7b)$$

$$\hat{\mathcal{P}}_z = \frac{\partial}{\partial z}. \quad (7c)$$

Let us gather all these vector fields into a single vector $\hat{\mathcal{X}} = (\hat{\mathcal{J}}_A, \hat{\mathcal{J}}, \hat{\mathcal{P}})$. The generalised velocity vector $\hat{\mathbf{q}}$ of the Lagrangian reads

$$\hat{\mathbf{q}} = \boldsymbol{\eta} \cdot \hat{\mathcal{X}}(\mathbf{q}) \equiv \sum_{i=1}^9 \eta_i \hat{\mathcal{X}}_i(\mathbf{q}), \quad (8)$$

as required by Poincaré's formalism. The equations of motion of \mathbf{C}_A , \mathbf{C} , and \mathbf{r} given by Eq. (8) are made explicit in the following. To obtain the equations of motion, we need the structure constants c_{ij}^k defined as

$$[\hat{\mathcal{X}}_i, \hat{\mathcal{X}}_j] \equiv \hat{\mathcal{X}}_i \hat{\mathcal{X}}_j - \hat{\mathcal{X}}_j \hat{\mathcal{X}}_i = c_{ij}^k \hat{\mathcal{X}}_k.$$

A direct calculation shows that the non-vanishing commutators are

$$\begin{aligned} [\hat{\mathcal{J}}_{A,x}, \hat{\mathcal{J}}_{A,y}] &= -\hat{\mathcal{J}}_{A,z}, & [\hat{\mathcal{J}}_x, \hat{\mathcal{J}}_y] &= \hat{\mathcal{J}}_z, \\ [\hat{\mathcal{J}}_{A,y}, \hat{\mathcal{J}}_{A,z}] &= -\hat{\mathcal{J}}_{A,x}, & [\hat{\mathcal{J}}_y, \hat{\mathcal{J}}_z] &= \hat{\mathcal{J}}_x, \\ [\hat{\mathcal{J}}_{A,z}, \hat{\mathcal{J}}_{A,x}] &= -\hat{\mathcal{J}}_{A,y}, & [\hat{\mathcal{J}}_z, \hat{\mathcal{J}}_x] &= \hat{\mathcal{J}}_y. \end{aligned} \quad (9)$$

To get Poincaré's equation, we also need to evaluate $\hat{\mathcal{X}}_i(f)$. We have

$$\hat{\mathcal{J}}_A(f) = \mathbf{0}, \quad (10a)$$

$$\hat{\mathcal{J}}(f) = \mathbf{G}_B \times (\boldsymbol{\Omega} + \boldsymbol{\Omega}_A) - \hat{\mathcal{J}}(U_{AB}), \quad (10b)$$

$$\hat{\mathcal{P}}(f) = \mu \mathbf{V} \times \boldsymbol{\Omega}_A - \hat{\mathcal{P}}(U_{AB}). \quad (10c)$$

The equation (10a) results from the invariance by rotation of the problem. Indeed, the Lagrangian does not depend on \mathbf{C}_A . In Eq. (10c), $\mathbf{V} = \mathbf{v} + \boldsymbol{\Omega}_A \times \mathbf{r}$ represents the velocity relative to the barycentric frame expressed in the body-fixed frame of A . In Eq. (10b), we have introduced $\mathbf{G}_B = \mathbf{I}_B(\boldsymbol{\Omega} + \boldsymbol{\Omega}_A)$, the angular momentum of the body B . Similarly, we denote by $\mathbf{G}_A = \mathbf{I}_A \boldsymbol{\Omega}_A$ the angular momentum of the body A , and by $\mathbf{L} = \mu \mathbf{r} \times \mathbf{V}$, the orbital angular momentum. The last ingredients in Poincaré's equations are the partial derivatives of the kinetic energy with respect to $\boldsymbol{\eta}$ which are

$$\frac{\partial T}{\partial \boldsymbol{\Omega}_A} = \mathbf{G}_A + \mathbf{G}_B + \mathbf{L}, \quad \frac{\partial T}{\partial \boldsymbol{\Omega}} = \mathbf{G}_B, \quad \frac{\partial T}{\partial \mathbf{V}} = \mu \mathbf{V}. \quad (11)$$

Combining Eqs. (8), (9), (10), (11), and Poincaré's equations

$$\frac{d}{dt} \frac{\partial T}{\partial \eta_i} = \sum_{j,k} c_{ij}^k \eta_j \frac{\partial T}{\partial \eta_k} + \hat{\mathcal{X}}_i(f),$$

we get

$$\begin{cases} \dot{\mathbf{C}}_A = \boldsymbol{\Omega}_A \cdot \hat{\mathcal{J}}_A(\mathbf{C}_A), \\ \dot{\mathbf{C}} = (\boldsymbol{\Omega}_B - \boldsymbol{\Omega}_A) \cdot \hat{\mathcal{J}}(\mathbf{C}), \\ \dot{\mathbf{r}} = \mathbf{r} \times \boldsymbol{\Omega}_A + \mathbf{V}, \end{cases} \quad \begin{cases} \dot{\mathbf{G}}_A = \mathbf{G}_A \times \boldsymbol{\Omega}_A + \mathbf{T}_A, \\ \dot{\mathbf{G}}_B = \mathbf{G}_B \times \boldsymbol{\Omega}_A + \mathbf{T}_B, \\ \dot{\mathbf{V}} = \mathbf{V} \times \boldsymbol{\Omega}_A + \frac{1}{\mu} \mathbf{F}, \end{cases} \quad (12)$$

with

$$\mathbf{F} = -\hat{\mathcal{P}}(U_{AB}), \quad \mathbf{T} = -\hat{\mathcal{L}}(U_{AB}), \quad \mathbf{T}_B = -\hat{\mathcal{J}}(U_{AB}), \quad \mathbf{T}_A = -\mathbf{T} - \mathbf{T}_B.$$

The rotation vectors are given by

$$\boldsymbol{\Omega}_A = \mathbf{I}_A^{-1} \mathbf{G}_A, \quad \boldsymbol{\Omega}_B = \mathbf{I}_B^{-1} \mathbf{G}_B,$$

and the vector field $\hat{\mathcal{L}}$ is defined as

$$\hat{\mathcal{L}}_x = y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y}, \quad (13a)$$

$$\hat{\mathcal{L}}_y = z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z}, \quad (13b)$$

$$\hat{\mathcal{L}}_z = x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x}. \quad (13c)$$

The equations (12) are strictly equivalent to [Maciejewski's](#) equations of motion, but they are explicitly written in terms of the vector fields $\hat{\mathcal{P}}$, $\hat{\mathcal{L}}$, $\hat{\mathcal{J}}_A$ and $\hat{\mathcal{J}}$. As shown in the subsequent section, textbooks on quantum theory of angular momentum allow to evaluate these vector fields applied to U_{AB} very efficiently.

2.2 Ladder operators, force and torques

We introduce a complex coordinate system $(\mathbf{e}_+, \mathbf{e}_0, \mathbf{e}_-)$ in which coordinates of any vector \mathbf{a} are denoted (a_+, a_0, a_-) and are related to the usual Cartesian coordinates (a_x, a_y, a_z) by

$$a_+ = -\frac{1}{\sqrt{2}}(a_x + ia_y) \quad a_0 = a_z \quad a_- = \frac{1}{\sqrt{2}}(a_x - ia_y).$$

We apply the same rule for vector fields $\hat{\mathcal{A}} = (\hat{\mathcal{A}}_x, \hat{\mathcal{A}}_y, \hat{\mathcal{A}}_z)$. We get

$$\hat{\mathcal{A}}_+ = -\frac{1}{\sqrt{2}}(\hat{\mathcal{A}}_x + i\hat{\mathcal{A}}_y) \quad \hat{\mathcal{A}}_0 = \hat{\mathcal{A}}_z \quad \hat{\mathcal{A}}_- = \frac{1}{\sqrt{2}}(\hat{\mathcal{A}}_x - i\hat{\mathcal{A}}_y).$$

The vector fields defined in Eqs. (5), (6), (7), and (13) are related to ladder operators commonly used in quantum theory of angular momentum. With the notation of [Varshalovich et al \(1988\)](#), we have

$$\hat{\mathcal{J}}_{A,\nu} \equiv i\hat{J}'^\nu(\alpha_A, \beta_A, \gamma_A), \quad \hat{\mathcal{J}}_\nu \equiv i\hat{J}_\nu(\alpha, \beta, \gamma), \quad \hat{\mathcal{P}}_\nu \equiv \nabla_\nu, \quad \hat{\mathcal{L}}_\nu \equiv i\hat{L}_\nu,$$

for $\nu \in \{+, 0, -\}$. \hat{J}'^ν and \hat{J}_ν are the components of the spin operator expressed in the rotated frame and in the initial frame, respectively. ∇_ν is the usual gradient operator and \hat{L}_ν is the orbital angular momentum operator. Given these equivalences, we deduce the following relations ([Varshalovich et al, 1988](#))

$$\hat{\mathcal{P}}_\nu \left(\frac{Y_{l,m}(\mathbf{r})}{r^{l+1}} \right) = \begin{cases} -\sqrt{(l+m+1)(l-m+1)} \frac{Y_{l+1,m}(\mathbf{r})}{r^{l+2}} & \nu = 0 \\ -\sqrt{\frac{(l \pm m + 1)(l \pm m + 2)}{2}} \frac{Y_{l+1,m \pm 1}(\mathbf{r})}{r^{l+2}} & \nu = \pm 1 \end{cases} \quad (14a)$$

and

$$\hat{\mathcal{L}}_\nu \left(\frac{Y_{l,m}(\mathbf{r})}{r^{l+1}} \right) = i \begin{cases} m \frac{Y_{l,m}(\mathbf{r})}{r^{l+1}} & \nu = 0 \\ \mp \sqrt{\frac{l(l+1) - m(m \pm 1)}{2}} \frac{Y_{l,m \pm 1}(\mathbf{r})}{r^{l+1}} & \nu = \pm 1 \end{cases}, \quad (14b)$$

for the orbital part (embedded in the spherical harmonics), while spin operators act on Wigner D-matrices according to

$$\hat{\mathcal{J}}_\nu D_{m,m'}^l(\mathbf{C}) = i \begin{cases} -m D_{m,m'}^l(\mathbf{C}), & \nu = 0, \\ \pm \sqrt{\frac{l(l+1) - m(m \mp 1)}{2}} D_{m \mp 1, m'}^l(\mathbf{C}) & \nu = \pm 1 \end{cases} \quad (15a)$$

and

$$\hat{\mathcal{J}}_{A,\nu} D_{m,m'}^l(\mathbf{C}_A) = i \begin{cases} -m' D_{m,m'}^l(\mathbf{C}_A), & \nu = 0, \\ \pm \sqrt{\frac{l(l+1) - m'(m' \pm 1)}{2}} D_{m, m' \pm 1}^l(\mathbf{C}_A) & \nu = \pm 1 \end{cases}. \quad (15b)$$

Forces and torques are then computed as follows. Let us define the constant terms $u_{l_2, m_2}^{l_1, m_1}$ as

$$u_{l_2, m_2}^{l_1, m_1} = -GM_A M_B R_A^{l_1} R_B^{l_2} Z_{l_1, m_1}^A (-1)^{l_2} \gamma_{l_2, m_2}^{l_1, m_1}. \quad (16)$$

The potential (Eq. 1) reads

$$U_{AB} = \sum_{l_1=0}^{\infty} \sum_{m_1=-l_1}^{l_1} \sum_{l_2=0}^{\infty} \sum_{m_2=-l_2}^{l_2} u_{l_2, m_2}^{l_1, m_1} Z_{l_2, m_2}^B(\mathbf{C}) \frac{Y_{l_1+l_2, m_1+m_2}(\mathbf{r})}{r^{l_1+l_2+1}},$$

and thus,

$$\begin{aligned} \mathbf{F} &= - \sum_{l_1=0}^{\infty} \sum_{m_1=-l_1}^{l_1} \sum_{l_2=0}^{\infty} \sum_{m_2=-l_2}^{l_2} u_{l_2, m_2}^{l_1, m_1} Z_{l_2, m_2}^B(\mathbf{C}) \hat{\mathcal{P}} \left(\frac{Y_{l_1+l_2, m_1+m_2}(\mathbf{r})}{r^{l_1+l_2+1}} \right), \\ \mathbf{T} &= - \sum_{l_1=0}^{\infty} \sum_{m_1=-l_1}^{l_1} \sum_{l_2=0}^{\infty} \sum_{m_2=-l_2}^{l_2} u_{l_2, m_2}^{l_1, m_1} Z_{l_2, m_2}^B(\mathbf{C}) \hat{\mathcal{L}} \left(\frac{Y_{l_1+l_2, m_1+m_2}(\mathbf{r})}{r^{l_1+l_2+1}} \right), \\ \mathbf{T}_B &= - \sum_{l_1=0}^{\infty} \sum_{m_1=-l_1}^{l_1} \sum_{l_2=0}^{\infty} \sum_{m_2=-l_2}^{l_2} u_{l_2, m_2}^{l_1, m_1} \hat{\mathcal{J}} \left(Z_{l_2, m_2}^B(\mathbf{C}) \right) \frac{Y_{l_1+l_2, m_1+m_2}(\mathbf{r})}{r^{l_1+l_2+1}}, \\ \mathbf{T}_A &= -\mathbf{T} - \mathbf{T}_B, \end{aligned}$$

with

$$Z_{l_2, m_2}^B(\mathbf{C}) = \sum_{m_2'=-l_2}^{l_2} D_{m_2, m_2'}^{l_2}(\mathbf{C}) Z_{l_2, m_2'}^B,$$

and

$$\hat{\mathcal{J}} \left(Z_{l_2, m_2}^B(\mathbf{C}) \right) = \sum_{m_2'=-l_2}^{l_2} \hat{\mathcal{J}} \left(D_{m_2, m_2'}^{l_2}(\mathbf{C}) \right) Z_{l_2, m_2'}^B.$$

The simplicity of the formulae (14) and (15) associated to the evaluation of the ladder operators makes the calculation of the components of \mathbf{F} , \mathbf{T} , and \mathbf{T}_B very efficient and easy to implement.

2.3 Spherical harmonics

Naturally, spherical harmonics are not computed from their definition Eq. (2) but from recurrence formulae which can also be found in many textbooks. Let $\mathbf{u} \equiv \mathbf{r}/\|\mathbf{r}\| = (u_+, u_0, u_-)$ be the unit vector along \mathbf{r} . By definition $Y_{l,m}(\mathbf{r}) \equiv Y_{l,m}(\mathbf{u})$. The initialisation is

$$Y_{0,0}(\mathbf{r}) = 1, \quad Y_{1,0}(\mathbf{r}) = u_0, \quad Y_{1,1}(\mathbf{r}) = u_+, \quad (17a)$$

and the recurrence equations are

$$lY_{l,0}(\mathbf{r}) = (2l-1)u_0Y_{l-1,0}(\mathbf{r}) - (l-1)Y_{l-2,0}(\mathbf{r}), \quad (17b)$$

$$\sqrt{l+m}Y_{l,m}(\mathbf{r}) = \sqrt{l-m}u_0Y_{l-1,m}(\mathbf{r}) + \sqrt{2(l+m-1)}u_+Y_{l-1,m-1}(\mathbf{r}). \quad (17c)$$

Spherical harmonics with negative order m are deduced from the symmetry relation

$$Y_{l,m}(\mathbf{r}) = (-1)^m \bar{Y}_{l,-m}(\mathbf{r}). \quad (17d)$$

As it can be noticed, in the set of equations (17), there is not any trigonometric functions.

2.4 Cayley-Klein parameters

To avoid singularities, rotations – involved in Wigner D-matrices – are parametrised by Cayley-Klein complex parameters a and b (e.g., [Varshalovich et al, 1988](#)) instead of the more common Euler 3-1-3 angles (ψ, θ, ϕ) or the Euler 3-2-3 angles (α, β, γ) . Cayley-Klein parameters are equivalent to quaternions, but are more adapted to the formalism used in this work. For the sake of completeness, we here summarise a few properties of these parameters ([Varshalovich et al, 1988](#)). They and Euler 3-1-3 angles are related to each other by

$$a = \cos \frac{\theta}{2} e^{-i\frac{\psi+\phi}{2}} = D_{\frac{1}{2}, \frac{1}{2}}^{\frac{1}{2}}(\mathbf{C}), \quad b = -i \sin \frac{\theta}{2} e^{i\frac{\psi-\phi}{2}} = D_{-\frac{1}{2}, \frac{1}{2}}^{\frac{1}{2}}(\mathbf{C}), \quad (18a)$$

and reciprocally,

$$\psi = \arg(i\bar{a}b), \quad \theta = \cos^{-1}(|a|^2 - |b|^2), \quad \phi = -\arg(iab). \quad (18b)$$

The expression of the Cartesian coordinates of a rotated vector, e.g. $\mathbf{r} = \mathbf{C}\mathbf{r}'$, expressed in terms of Cayley-Klein parameters is

$$r_0 = (|a|^2 - |b|^2)r'_0 + 2\sqrt{2}\Re(ab\bar{r}'_+) \quad r_+ = -\sqrt{2}\bar{a}br'_0 + \bar{a}^2r'_+ - b^2\bar{r}'_+,$$

where $\Re(z)$ means the real part of z . The inverse rotation is obtained by the substitution $(a, b) \rightarrow (\bar{a}, -b)$. The product of two rotations $\mathbf{C}(a, b) = \mathbf{C}(a_1, b_1)\mathbf{C}(a_2, b_2)$ is given by

$$a = a_1a_2 - \bar{b}_1b_2, \quad b = b_1a_2 + \bar{a}_1b_2.$$

Finally, the equivalent of [Maciejewski](#)'s equations of motion of \mathbf{C} and \mathbf{C}_A (Eq. 12) in terms of Cayley-Klein parameters (a, b) are deduced from their expressions in terms of Wigner D-matrix (Eq. 18a). Indeed, for all (l, m, m') , we have

$$\begin{cases} \dot{D}_{m,m'}^l(\mathbf{C}) &= (\boldsymbol{\Omega}_B - \boldsymbol{\Omega}_A) \cdot \hat{\mathcal{J}}(D_{m,m'}^l(\mathbf{C})) \\ \dot{D}_{m,m'}^l(\mathbf{C}_A) &= \boldsymbol{\Omega}_A \cdot \hat{\mathcal{J}}_A(D_{m,m'}^l(\mathbf{C}_A)) \end{cases} .$$

Taking the particular values of (l, m, m') corresponding to the definitions of a and b (Eq. 18a), we get

$$\begin{cases} \dot{a} = -\frac{i}{2} \left(a(\Omega_{B,0} - \Omega_{A,0}) - \sqrt{2}b(\bar{\Omega}_{B,+} - \bar{\Omega}_{A,+}) \right) \\ \dot{b} = +\frac{i}{2} \left(b(\Omega_{B,0} - \Omega_{A,0}) + \sqrt{2}a(\Omega_{B,+} - \Omega_{A,+}) \right) \end{cases} \quad (19a)$$

and

$$\begin{cases} \dot{a}_A = -\frac{i}{2} \left(a_A \Omega_{A,0} + \sqrt{2}\bar{b}_A \Omega_{A,+} \right) \\ \dot{b}_A = -\frac{i}{2} \left(b_A \Omega_{A,0} - \sqrt{2}\bar{a}_A \Omega_{A,+} \right) \end{cases} . \quad (19b)$$

2.5 Wigner D-matrices

Like for spherical harmonics, the computation of Wigner D-matrices is performed recursively. Here, we need these matrices with integer l . The recurrence is initialised with

$$D_{0,0}^0 = 1, \quad D_{m,m'}^1 = \begin{bmatrix} 1 & 0 & -1 \\ a^2 & -\sqrt{2}a\bar{b} & (\bar{b})^2 \\ \sqrt{2}ab & |a|^2 - |b|^2 & -\sqrt{2}\bar{a}\bar{b} \\ b^2 & \sqrt{2}\bar{a}b & (\bar{a})^2 \end{bmatrix} \begin{matrix} \hat{\Gamma} \\ \hat{\Theta} \\ \hat{\Upsilon} \\ \hat{\Xi} \end{matrix} .$$

For $m \geq 0$, Wigner D-matrices of degree $l \geq 2$ are given by (see [Gimbutas and Greengard, 2009](#))

$$D_{m,m'}^l = c_{m,m'}^{l,-} D_{1,1}^1 D_{m-1,m'-1}^{l-1} + c_{m,m'}^{l,0} D_{1,0}^1 D_{m-1,m'}^{l-1} + c_{m,m'}^{l,+} D_{1,-1}^1 D_{m-1,m'+1}^{l-1} \quad (20a)$$

with coefficients

$$\begin{cases} c_{m,m'}^{l,-} = \sqrt{\frac{(l+m')(l+m'-1)}{(l+m)(l+m-1)}}, \\ c_{m,m'}^{l,0} = \sqrt{\frac{2(l+m')(l-m')}{(l+m)(l+m-1)}}, \\ c_{m,m'}^{l,+} = \sqrt{\frac{(l-m')(l-m'-1)}{(l+m)(l+m-1)}}. \end{cases} \quad (20b)$$

If $|m'+\nu|$, with $\nu \in \{-1, 0, 1\}$, is strictly greater than $l-1$, then $D_{m-1, m'+\nu}^{l-1}$ should be discarded and replaced by zero in the left-hand side of Eq. (20a). Elements of Wigner D-matrices with negative index m are deduced from the symmetry relation

$$D_{m, m'}^l = (-1)^{m-m'} \bar{D}_{-m, -m'}^l .$$

3 Numerical implementation

The algorithm presented above has been implemented in C++. To fasten the integration, recurrence coefficients in Eqs. (17b), (17c) and (20b), the constant terms $u_{l_2, m_2}^{l_1, m_1}$ (Eq. 16), as well as the constant values in the evaluation of the ladder operators (Eqs. 14 and 15), are all calculated and stored before the effective integration of the equations of motion (12). To be homogeneous with Hou et al (2016), a truncation of the mutual potential at order n means that $l_1 + l_2 \leq n$ (l_1 runs from 0 to n and l_2 from 0 to $n - l_1$) in the expression of U_{AB} , \mathbf{F} , \mathbf{T} , and \mathbf{T}_B (Sect. 2.2).

In order to compare the efficiency of the algorithm with that presented by Hou et al (2016), we integrate the binary asteroid system 1999 KW4 (66391). The polyhedral shapes of the two components are retrieved at the URL (<http://echo.jpl.nasa.gov/asteroids/shapes/>). Inertia integrals

$$T^{l, m, n} = \int \rho x^l y^m z^n dx dy dz,$$

with ρ the (constant) density of the considered body and where the integration is done over its whole volume, are computed using the formulae of Hou et al (2016). In particular, inertia integrals of degree $l + m + n = 0$ and 1 are used to compute the coordinates of the barycenter which are then subtracted to those of the polyhedron vertexes. This procedure ensures that the body-fixed frame of each component is well centred on the barycenter as required by the equations of motion (Eq. 12). The generalised products of inertia are then converted into Stokes coefficients following Tricarico (2008) but with a different normalisation factor¹. Stokes coefficients can alternatively be directly computed from the polyhedron model using the algorithm of Werner (1997). We assume the following masses: $M_A = 2.355 \times 10^{12}$ kg and $M_B = 0.135 \times 10^{12}$ kg, which implies that the densities of the two components are 1970.2 kg.m^{-3} and 2810.5 kg.m^{-3} , respectively.

The initial conditions are the same as in (Hou et al, 2016): the orbital elements are

$$a = 2.5405 \text{ km}, \quad e = 0.01, \quad i = \Omega = \omega = 0^\circ, \quad M = 180^\circ .$$

¹ If we denote by $\{C_{lm}^{(T)}, S_{lm}^{(T)}\}$ Stokes coefficients defined in (Tricarico, 2008, Eqs. 14, 15), those of the present paper are given by $\{C_{lm}, S_{lm}\} = N_{lm} \{C_{lm}^{(T)}, S_{lm}^{(T)}\}$ with

$$N_{lm} = \frac{2}{1 + \delta_{m,0}} \frac{(l-m)!}{(l+m)!} .$$




Fig. 1 Orbital evolution of the 1999 KW4 binary asteroid system over 10 000 h time span obtained with an RKF7(8) integrator with a fixed time step of 200 s. The mutual potential is truncated at order 6. a is the semi-major axis, e the eccentricity, and i the inclination relative to the initial orbital plane.

The initial angular speeds of the two components in their respective body-fixed frame are

$$\boldsymbol{\Omega}_A = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} 315.4^\circ \cdot \text{day}^{-1}, \quad \boldsymbol{\Omega}_B = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} 495^\circ \cdot \text{day}^{-1}.$$

The 3-1-3 Euler angles defining the initial orientations of the body-fixed reference frames relative to the initial orbit plane are

$$\psi_A = 27.04^\circ, \quad \theta_A = 10^\circ, \quad \phi_A = -83.93^\circ \quad \psi_B = 0^\circ, \quad \theta_B = 0^\circ, \quad \phi_B = 180^\circ.$$

In order to make the most objective comparison with [Hou et al \(2016\)](#), we use the same RKF7(8) integrator with the same fixed time step of 200 seconds. Tests are run on a single processor with a CPU frequency of 2.3 GHz.

Figure 1 represents the orbital evolution of the system with respect to the initial orbital plane over 10 000h. The mutual potential is truncated at order 6. As observed by [Hou et al \(2016\)](#), the energy stored in the rotational motion is able to strongly influence the orbit by forcing its eccentricity and inclination to oscillate in the intervals $0 \leq e \leq 0.035$ and $0 \leq i \leq 16^\circ$, respectively. The total energy of the system is 8.7678×10^{10} J. Numerically, its variation is mainly due to the dissipative effect of the RKF integrator. Over the 10 000 h of integration time span, the total variation of the mechanical energy is only -0.3870 J, which is perfectly consistent with the -0.3851 J obtained by [Hou et al \(2016\)](#).

A comparison of the integration times between our approach and the algorithm of [Hou et al \(2016\)](#) is presented in Table 1. The system is integrated over 20 000 h

Table 1 Computation time for a 200 h integration of the orbit using [Hou et al](#)'s algorithm (first row) and our approach (second row).

Order	2	3	4	5	6	7	8	9
t^a (sec)	0.2	0.7	2.1	6.6	17.3	42.6	99.3	206.0
t^b (sec)	0.2	0.3	0.4	0.6	1.0	1.7	2.5	3.1

t^a computation time using [Hou et al](#)'s algorithm on a single 2.9 GHz CPU frequency processor.

t^b computation time using the present algorithm on a single 2.3 GHz CPU frequency processor.

with a fixed time step of 200 s, and then the computation time is divided by 100 for a direct comparison with [Hou et al \(2016\)](#) who did the integration of the orbit over 200 h. Absolute CPU times highly depend on the compiler and the optimisation options. They should thus be compared with care. Nevertheless, both methods spend the same amount of computation time when the potential is truncated at the lowest order $n = 2$. We can thus assume that the other columns of [Tab. 1](#) objectively reflect the efficiency of the algorithms. Moreover, as the expansion order increases, the computation time growth differs significantly between the two methods. Ours proves to be much faster. At the 6th order of truncation, we already gain a factor 17 in speed, and at order 9, our approach only needs 3 seconds instead of the 3.4 minutes required by the algorithm of [Hou et al \(2016\)](#).

4 Conclusion

The decomposition of the mutual potential of two rigid bodies with arbitrary shape into spherical harmonics has sometimes been discarded because of the misconception that this formalism would involve trigonometric functions. However spherical harmonics do have expressions in terms of Cartesian coordinates. Moreover, not only recurrence relations allow to evaluate them efficiently, but the force and the orbital torque are easily computed by application of the gradient and the angular momentum operators, respectively. By consequence, if we restrain ourselves to the orbital part of the equations of motion only, our approach must be as efficient as a polynomial decomposition of the potential in Cartesian coordinates.

Nevertheless, numerical tests show that our algorithm is faster. The reason resides in the way rotations are handled. Here, we use the irreducible representation of the group $SO(3)$ acting on the set of functions defined on the sphere, viz. Wigner D-matrices. These matrices have two advantages: they can be efficiently evaluated with recurrence relations and the torque is simply obtained by application of the spin operator. Combining spherical harmonics and Wigner D-matrices, we obtain an algorithm for which forces and torques are calculated as rapidly as the mutual potential.

In this work, in order to avoid singularities, rotations are parametrised using Cayley-Klein's formalism whose parameters are strictly equivalent to quaternions. We make this choice because it is well adapted to the computation of Wigner D-matrices. Because this representation is not common in the celestial mechanics field, we briefly summarise the relation between these parameters and the standard 3-1-3 Euler angles. More properties can be found in textbooks on quantum theory of angular momentum (e.g., [Varshalovich et al, 1988](#)).

As a concluding remark, not only the present formalism allows to design a fast integrator for the full two rigid body problem, but it also reveals the underlying structure which is controlled by the group of rotations. This structure is often hidden in other approaches. We expect that this will allow to draw a more comprehensive view of the analytical problem.

References

- Borderies N (1978) Mutual gravitational potential of N solid bodies. *Celestial Mechanics* 18:295–307, DOI 10.1007/BF01230170
- Boué G, Correia ACM, Laskar J (2016) Complete spin and orbital evolution of close-in bodies using a Maxwell viscoelastic rheology. *Celestial Mechanics and Dynamical Astronomy* DOI 10.1007/s10569-016-9708-x
- Compère A, Lemaître A (2014) The two-body interaction potential in the STF tensor formalism: an application to binary asteroids. *Celestial Mechanics and Dynamical Astronomy* 119:313–330, DOI 10.1007/s10569-014-9568-1
- Fahnestock EG, Scheeres DJ (2006) Simulation of the full two rigid body problem using polyhedral mutual potential and potential derivatives approach. *Celestial Mechanics and Dynamical Astronomy* 96:317–339, DOI 10.1007/s10569-006-9045-6
- Gimbutas Z, Greengard L (2009) A fast and stable method for rotating spherical harmonic expansions. *Journal of Computational Physics* 228:5621 – 5627
- Hartmann T, Soffel MH, Kioustelidis T (1994) On the use of STF-tensors in celestial mechanics. *Celestial Mechanics and Dynamical Astronomy* 60:139–159, DOI 10.1007/BF00693097
- Hou X, Scheeres DJ, Xin X (2016) Mutual Potential between Two Rigid Bodies with Arbitrary Shapes and Mass Distributions. *Celestial Mechanics and Dynamical Astronomy* pp 1–27, DOI 10.1007/s10569-016-9731-y
- Maciejewski AJ (1995) Reduction, Relative Equilibria and Potential in the Two Rigid Bodies Problem. *Celestial Mechanics and Dynamical Astronomy* 63:1–28, DOI 10.1007/BF00691912
- Mathis S, Le Poncin-Lafitte C (2009) Tidal dynamics of extended bodies in planetary systems and multiple stars. *Astron. Astrophys.* 497:889–910, DOI 10.1051/0004-6361/20079054
- Paul MK (1988) An Expansion in Power Series of Mutual Potential for Gravitating Bodies with Finite Sizes. *Celestial Mechanics* 44:49–59, DOI 10.1007/BF01230706
- Poincaré H (1901) Sur une forme nouvelle des équations de la mécanique. *Comptes rendus de l'Académie des Sciences* 132:369–371
- Tricarico P (2008) Figure figure interaction between bodies having arbitrary shapes and mass distributions: a power series expansion approach. *Celestial Mechanics and Dynamical Astronomy* 100:319–330, DOI 10.1007/s10569-008-9128-7, [0711.2078](#)
- Varshalovich D, Moskalev A, Khersonskii V (1988) *Quantum Theory of Angular Momentum*. World Scientific
- Werner RA (1997) Spherical harmonic coefficients for the potential of a constant-density polyhedron. *Computers and Geosciences* 23:1071–1077, DOI 10.1016/S0098-3004(97)00110-6

Werner RA, Scheeres DJ (2005) Mutual Potential of Homogeneous Polyhedra. *Celestial Mechanics and Dynamical Astronomy* 91:337–349, DOI 10.1007/s10569-004-4621-0

Wigner EP (1959) *Group theory and its application to the quantum mechanics of atomic spectra*. Academic Press, New York