



Ecology of termites from the genus *Nasutitermes* (Termitidae: Nasutitermitinae) and potential for science-based development of sustainable pest management programs

Isabelle Boulogne, Reginaldo Constantino, Nadine Amusant, Michaël Falkowski, Alice M. S Rodrigues, Emeline Houël

► To cite this version:

Isabelle Boulogne, Reginaldo Constantino, Nadine Amusant, Michaël Falkowski, Alice M. S Rodrigues, et al.. Ecology of termites from the genus *Nasutitermes* (Termitidae: Nasutitermitinae) and potential for science-based development of sustainable pest management programs. *Journal of Pest Science*, 2017, 90 (1), pp.19 - 37. 10.1007/s10340-016-0796-x . hal-01467356

HAL Id: hal-01467356

<https://hal.sorbonne-universite.fr/hal-01467356>

Submitted on 14 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Ecology of termites from the genus *Nasutitermes* (Termitidae: Nasutitermitinae) and potential for science-based development of sustainable pest management programs

Isabelle Boulogne^{a,b,c*}, Reginaldo Constantino^d, Nadine Amusant^e, Michaël Falkowski^b, Alice M.S. Rodrigues^f, Emeline Houël^b

^a Université des Antilles, Campus de Fouillole, 97157 Pointe-à-Pitre Cedex, Guadeloupe, FRANCE.

^b CNRS, UMR EcoFoG (AgroParisTech, CIRAD, INRA, UA, UG), Institut Pasteur de la Guyane, 23 Avenue Pasteur, BP6010, 97306 Cayenne Cedex, French Guiana, FRANCE.

^c Université Pierre et Marie Curie, UMR 7618, Institut d'Ecologie et des Sciences de l'Environnement de Paris (iEES-Paris), Département d'Ecologie Sensorielle, 75252 Paris Cedex 05, FRANCE.

^d Departamento de Zoologia, Universidade de Brasília, 70910-970 Brasília, DF, BRASIL.

^e CIRAD, UMR EcoFoG (AgroParisTech, CNRS, INRA, UA, UG), BP701, 97310 Kourou Cedex, French Guiana FRANCE.

^f Sorbonne Universités, Université Pierre et Marie Curie, CNRS, Laboratoire de Biodiversité et Biotechnologies Microbiennes (LBBM), Observatoire Océanologique, 66650 Banyuls/Mer, FRANCE.

* Corresponding author: Isabelle BOULOGNE.

Current professional adress: Université Pierre et Marie Curie, UMR 7618, Institut d'Ecologie et des Sciences de l'Environnement de Paris (iEES-Paris), Département d'Ecologie Sensorielle, 75252 Paris Cedex 05.

E-mail address: isabelle.boulogne@univ-ag.fr or isabelle.boulogne@upmc.fr

Key Message

- We reviewed the taxonomic status of *Nasutitermes*, which is currently the most diverse termite genus, has a particularly complex taxonomic history, and includes some major pest species.
- We performed a comparative literature analysis, especially concerning the type-species *N. corniger*, while filtering available information concerning its biology through the prism of termite control.
- We critically examined the existing management of *N. corniger* (synthetic chemical products, botanical insecticides towards various targets, and biological control) to lead to the development of innovative management tools and strategies.

Author Contribution Statement

IB conceived the first draft. RC wrote the taxonomic history part and corrected biological part. EH, AR and NA corrected and greatly contributed to the pest management part. MF made all the drawings of *N. corniger*'s colony. All authors read, corrected and approved the manuscript.

Abstract

The genus *Nasutitermes* is among the most abundant wood-feeding Termitidae and an extremely diverse and heterogeneous group in terms of its biogeography and morphology. Despite the major role of several *Nasutitermes* species as structural pests, the phylogenetic status of this genus is still unclear, alongside with a confused taxonomy and species identification remaining difficult. The first aim of this review was thus to gather and discuss studies concerning the taxonomic status of the genus *Nasutitermes*, in order to clarify this crucial point. Then, our goal was to gain new insights into the management of *N. corniger*, considered to be the most economically detrimental pest of this genus in South America and a *Nasutitermes* model species, while filtering available information concerning its biology through the prism of termite control, as well as critically examine the existing methods. We indeed strongly believe that increasing our knowledge of this species' biological strategies is the key to progress in the challenging question of their sustainable management.

Keywords: Taxonomic history, *Nasutitermes corniger*, Termitidae, sustainable management, IPM, antimicrobial and insecticidal botanical extracts.

1. Introduction

The species causing the most considerable damage to tropical agriculture belong to four of the eight subfamilies of higher termites (Termitidae), and Nasutitermitinae is one of them (Rouland-Lefèvre 2011; Krishna et al. 2013). The genus *Nasutitermes* is among the most abundant wood-feeding Nasutitermitinae in the tropics and several *Nasutitermes* species are important structural pests (Constantino 2002; Fontes and Milano 2002). This genus comprises a diverse and heterogeneous group of species (Miura et al, 2000). The distributional patterns of *Nasutitermes* species were probably shaped by environmental conditions and historical factors, such as continental drift, orogeny, and Quaternary climatic variations (Miura et al, 2000).

As currently defined, *Nasutitermes* does not comprise a monophyletic group of species (Eggleton 2001; Inward et al. 2007; Roy et al. 2014), its taxonomy is confused, and accurate species identification is very difficult. Many nasute species that do not belong to the *Nasutitermes* clade remain nested within *Nasutitermes*, generating taxonomic confusion, while misidentifications and nomenclatural errors are frequent (Gush et al. 1985; Traniello et al. 1985; Constantino 2000; 2002; Scheffrahn et al. 2002; Scheffrahn et al. 2005a; Scheffrahn et al. 2005b). A comprehensive taxonomic revision of the genus is highly necessary, but has never been attempted due to the enormous difficulties involved.

However, despite this heterogeneity, some aspects, such as nest building (Thorne et al. 1996; Fuller and Postava-Davignon 2014), soldiers' defense strategies (Prestwich 1979) and hindgut microbiome role in symbiosis (Brune 2014), are well known for several *Nasutitermes* species. Yet, this information could be more clearly highlighted and discussed in the perspective of pest management to exploit at its most and face the growing importance of *Nasutitermes* pest species. These termites live in a large range of habitats in urban, anthropized, disturbed and natural environments, in dry as well as in moist conditions and at

elevations of up to 1000 m (Scheffrahn et al. 2005a; Bustamante and Martius 1998; Dunn and Messier 1999, Vasconcellos and Moura 2010, Mello et al. 2014, Lima et al. 2013). This versatility, alongside with changes in urban soil occupation in the last few decades, leads to an increasing infestation in large buildings which are full of structural voids, narrow fissures and wood (Fontes and Milano 2002).

In this paper, we thus gathered and commented studies on taxonomic history and current status of genus *Nasutitermes*, in order to clarify this crucial point. Indeed, taxonomy provides an essential framework, as no effective and sustainable management can be set up without precisely identifying and describing the intended species.

We also aimed to gain new insights into the biology of *N. corniger* as a *Nasutitermes* model species, in the perspective of improving termite control. Better insights into these aspects would indeed contribute to the development of more adapted and efficient control methods, as highlighted by Scharf (2015) in a recent review.

Eventually, we critically examined the existing management methods against *N. corniger* and lastly underlined the fact that termites are also beneficial, playing a key role in the wood decomposition process.

2. Taxonomic history of genus *Nasutitermes* Dudley, 1890

Nasutitermes has a complex taxonomic history and many of the species currently included in this genus have been classified in other termite genera, while several species previously included in *Nasutitermes* have been transferred to other genera.

Our current taxonomic system begins with Linnaeus (1758), who described a single termite genus, *Termes*, which means termite in Latin. All termite species were classified in genus *Termes* for nearly a century after Linnaeus. In 1781 the British naturalist Henry Smeathman, described *Nasutitermes arborum* (as *Termes arborum*) from Sierra Leone,

Africa, which was the first species of *Nasutitermes* to be formally named. Smeathman (1781) also presented information on its biology, nests, and damage to houses.

Termite classification started to change in the second half of the 19th century, when the German entomologist H.A. Hagen was the main authority on termite taxonomy. Hagen's (1858) monograph on world termites listed seven species of *Nasutitermes*, all of them included in the subgenus *Termes* (*Eutermes*) Heer, 1849. *Eutermes* was defined based on wing venation and in Hagen scheme it included a heterogeneous group of species, currently classified in several distinct genera.

The concept of “*Eutermes*” was restricted by Brauer (1868) to include only the termite species with a nasute soldier. Müller (1873), Froggatt (1897), and Silvestri (1903) adopted Brauer's definition and also raised *Eutermes* to the status of a genus. During that period, “*Eutermes*” was equivalent to our current definition of the subfamily Nasutitermitinae, and included all known species of *Nasutitermes*.

After the discovery of many new termite species with nasute soldiers, Holmgren (1910, 1912) subdivided the genus *Eutermes* into several subgenera. In Holmgren's system, subgenus *Eutermes* (*Eutermes*) was approximately equivalent to our current definition of genus *Nasutitermes*.

The name *Nasutitermes* itself appears for the first time in a paper about termites of Panama (Dudley 1890). Because Dudley did not provide a formal taxonomic description and did not include any species in the new genus, several authors overlooked the new name. It is important to note that formal international rules of nomenclature did not exist in 1890 (the first edition of the Code was published in 1905). Despite its limitations, Dudley's description satisfies the rules of nomenclature established in the first edition of the Code, and is therefore an available name.

Banks (1918) revealed a major problem with the previous use of the name “*Eutermes*”: its type-species, *Eutermes debilis*, a fossil, was identified as a species of *Microcerotermes*, a genus with mandibulate soldiers not related to the nasutes. In the same paper, Banks adopted the name *Nasutitermes* Dudley, 1890 for the termites with nasute soldiers. The status of the name *Eutermes* is still considered uncertain and it has not been used for living termites. However, many species currently placed in *Nasutitermes* appeared in the literature under the name “*Eutermes*” for a couple of decades, including the important monographs published by Holmgren (1909, 1910, 1912).

The name *Nasutitermes* was widely adopted after 1918. Initially, it included all species with nasute soldiers (= subfamily Nasutitermitinae), and was later subdivided into several new subgenera and genera. Snyder's (1949) world catalog of termites lists about 400 species of nasute termites, classified into 25 different genera; among them, 190 species were listed under the genus *Nasutitermes*.

Today, *Nasutitermes* is the most diverse termite genus, with 254 valid species (Krishna et al. 2013, updated). It is present in all biogeographical regions, distributed as follows: Oriental 114, Neotropical 78, Papuan 20, Ethiopian 19, Australian 19, Palearctic 14, and Nearctic 01. As shown in Figure 1, new species of *Nasutitermes* have been discovered continually since the end of the 19th century. Variation in the description rate is related to the number of active termite taxonomists, which was larger during the first half of the 20th century. The total number of species is likely to increase in the future.

The species currently included in genus *Nasutitermes* do not form a monophyletic group (Inward et al. 2007). Some are quite distinct from the type species (*N. corniger*) and were included in *Nasutitermes* when this genus comprised most Nasutitermitinae. The reexamination of the status of these species has resulted in changes in the classification, with the description of new genera and reassignment of some species to other known genera of

Nasutitermitinae. For instance, several species previously included in *Nasutitermes* were transferred to *Cortaritermes* by Mathews (1977), Fontes (1998), and Cuezco et al. (2015). The name *Nasutitermes*, however, will always be associated with its type-species, *N. corniger*.

3. *Nasutitermes corniger* (Motschulsky, 1855) as a model species

Nasutitermes corniger is a neotropical species which occurs natively from southern Mexico to northern Argentina and in the Caribbean islands, and it is an invasive species in the Bahamas, Florida and New Guinea (Figure 2) (Scheffrahn et al. 2005a; Evans et al. 2013). In the latter area, it may have been introduced accidentally due to human transport (Scheffrahn et al. 2005b). It is widely distributed in the neotropics, with high adaptability to a wide range of habitats (Scheffrahn et al. 2002). *Nasutitermes corniger* is the most important pest species of this genus in the New World (Constantino 2002; Fontes and Milano 2002) and its biology is relatively well-known.

3.1 Biology, ecology, and behavior of *Nasutitermes corniger*

Apterous line: tasks and morphological polyethism

Nasutitermes corniger is a social insect that has colonies that can contain up to 900,000 individuals (Dunn and Messier 1999; Thorne 1984). A colony is composed of workers, soldiers, and imagoes. In Termitidae, this structure originates from a common first stage of development, from which an apterous and an imaginal concurrently emerge. The imaginal line leads to the reproductive caste. From the apterous line the worker caste appears, followed by soldiers (Figure 3) (Lima et al. 2013).

Workers are primarily responsible for foraging, constructing and repairing the nest; carrying eggs and nursing; and feeding and grooming immature stages, soldiers, and reproductives as well as participating in defensive activities with soldiers (Thorne 1984). Soldiers are a defensive caste responsible for guarding the colony (Traniello 1981; Gazal et al. 2012; Verma et al. 2009). They also organize foraging by exploring new wood sources (Traniello 1981) and by regulating foraging activity. During nest defense, soldiers squirt from their nasus a sticky, odoriferous, irritating secretion to distances of more than three times their body length (Traniello 1981; McMahan 1982). All sterile forms (workers and soldiers) are blind and lucifugous (Thorne 1983; Scheffrahn et al. 2005a; Verma et al. 2009). These termite castes are 4 to 6 mm long, pale and soft bodied (Figure 4); mature in one year; and can live up to 5 years (Verma et al. 2009). As is typical for the genus *Nasutitermes*, workers show sexual dimorphism, with females conspicuously larger than males (Lima et al. 2013; Scheffrahn et al. 2005a; Jones 1980). Soldiers represent between 5 and 20% of a colony (Dunn and Messier 1999; Thorne 1984). They are characterized by reduced mandibles and a modified cephalic projection (the nasus), giving its name to the genus (Thorne 1984). Both soldiers and workers are completely sterile, with vestigial reproductive organs.

In a recent study, Lima et al. (2013) studied caste polymorphism of *N. corniger* apterous line using morphometric and discriminant analysis. Based on the comparison of the length of the antenna, thorax and metatibia, they highlighted two morphological types in presoldiers and soldiers, and two lines of development for workers. These elements demonstrate that morphological studies of *N. corniger* still remain crucial to increase the knowledge of the caste system in this species. Coupled with analytical and statistical tools, they can still lead to new discoveries. In addition to the rigorous determination of morphological types, these elements contribute to improving the understanding of individuals' interactions inside the colony. Lima et al. (2013) emphasized the fact that

knowing whether the different morphological types perform different tasks would help in understanding the behavioral dynamics in termites' societies. Former studies have, for example, shown that the behavior of each caste is significantly different during nest repair and nest expansion (McMahan 1970). Moreover, molecular approaches also offer promise for deciphering cryptic aspects of caste biology, potentially leading to insights for new control technologies (Scharf et al., 2015).

Nymphal line: Description of the reproductive strategies

Imagoes are of three types: primary reproductives (queens and kings), alates, and adultoids (Thorne 1984). They have fully developed eyes and are winged and pigmented (Scheffrahn et al. 2005a; Verma et al. 2009; Thorne 1983). The primary reproductives are the imagoes that initiated the colony after the dispersal flight. Alates are produced seasonally and leave the nest after spending 5 to 8 months within the parental colony, while adultoids are replacement reproductives derived from alates that did not disperse (Thorne 1983).

Ergatoids are worker-like neotenic reproductives derived from immatures of the neuter line. They have two additional molts compared to worker's development. They are wingless, blind and are at the bottom of the fecundity hierarchy (Figure 3) (Thorne 1984).

Mature colonies of *N. corniger* release large numbers of alates during massive crepuscular nuptial flights (35% of the colony biomass and 5000 to 25000 individuals on average), usually after rain (Thorne 1983). Royal couples are formed, find moist places with wood food, shed their wings, and remain together throughout the life cycle of the colony. Queens have a physogastric abdomen (due to ovariole growth), are capable of laying approximately 3000 eggs per day, and can live up to 25 years (Scheffrahn et al. 2005a). The eggs are yellow to white and incubate for 50 to 60 days before hatching (Verma et al. 2009).

Colonies of *N. corniger* have the largest number of primary queens per colony and the highest proportion of polygynous colonies, as 20 to 25% of mature colonies are headed by multiple unrelated primary reproductives (Thorne 1984; Atkinson and Adams 1997; Hartke and Rosengaus 2013). The tolerance of several queens in polygynous nests was hypothesized to be due to the high individual cost of intensively laying at high rates, and this phenomenon is apparently inversely related to the mean weight of queens (Adams and Atkinson 2008). However, a study performed by Hartke and Rosengaus regarding the cost of pleometrosis compared with colony foundation by monogamous pairs demonstrated that this strategy is disadvantageous for the colony's fitness (Hartke and Rosengaus 2013). Thus, mature polygamous nests of *N. corniger* may not be established via pleometrosis, but may imply other phenomena, such as the coalescence of young monogamous colonies. In fact, polygyny in *N. corniger*, as in the case of other Isoptera and some Hymenoptera, is not accompanied by nepotism because no data showed that workers care for their mothers more than the other queens (Atkinson et al. 2008). Likewise, experimental young colonies easily merged under laboratory conditions (Hartke and Rosengaus 2013). These results highlight the fact that further studies may be useful to understand the exact process as well as the ecological advantages of this phenomenon of colony fusion. Alongside its feeding habits and the aspects related to nesting, these reproduction mechanisms may play a part in the species' success in colonizing space, thus making *N. corniger* a major urban pest (Mello et al. 2014).

Role of the nest: from structural protection to immunity enhancement

Nests of *N. corniger* (Figure 5) are constructed on trees, bases of trunks, in secluded places of buildings or, less often, on the open soil (Scheffrahn et al. 2005a). These nests are single or polycalic, from ovoid to conical in form, with dimensions ranging up to 90 cm in height and 40 cm in width (Dunn and Messier 1999; McMahan 1982; Levings and Adams

1984; Scheffrahn et al. 1990). The nests have three parts: a superficial area with a thin skin of carton, which can exclude predators and prevent desiccation and flooding conditions (McMahan 1982; Fuller and Postava-Davignon, 2014); a middle part consisting of interconnecting galleries and chambers made of hard and dense carton (McMahan 1982); and a queen chamber located near the center of the nest (Thorne 1980). Additionally, carton-covered tunnels radiate from the nest and provide protection to termites during foraging (Levings and Adams 1984). This particular organization represents a challenge in termite management, because nests are often distant from infestation sites and are so structurally sound, *Nasutitermes* control can thus often be difficult (Fontes and Milano 2002; Gazal et al. 2012).

The stercoral carton used by termite workers during foraging, constructing or repairing processes is a black substance mainly composed of masticated wood and feces, cemented with salivary secretions (Postava-Davignon 2010; Traniello 1981; McMahan 1982). In addition, some other chemical or biological substances can also be incorporated inside this material. For example, $\beta(1,3)$ -glucanase activity was recently discovered in the *N. corniger* nest carton due to the presence of the native tGNBP-2 protein. By cleaving and releasing pathogenic components, it participates in the colony's immunological defense system against infections (Bulmer et al. 2004; 2009). Also, preliminary results obtained on *N. acajutlae* highlighted that, in general, fewer bacteria and fungi were observed inside the nests compared to the soil and trails (Postava-Davignon 2010). Thus, *Nasutitermes* nests are effective structures that allow the insects to survive in their environment, and antimicrobial defense integrated in the structure can help termites to enhance their immunity. This perfectly exemplifies the richness of defense strategies among insect colonies. Based on this information, Bulmer et al. (2009) developed a glycomimetic molecule blocking tGNBP-2 using rational design, which increases infection and death rate of exposed termites and can be used as an alternative control method.

This work clearly demonstrated that understanding termite biology is key to their sustainable management, as well as the resolution of product engineering and development issues.

Pheromone biology: communication, recruitment, modulation

Chemical odor trails are produced by the sternal gland (Stuart 1963). This gland is located on the fifth abdominal sternite and is composed of modified epidermal cells (Traniello and Busher 1985). Soldiers' sternal glands are smaller, on average, than workers'. These sternal pheromones are implicated in several aspects of the colony life. When nest defense is necessary, soldiers release a secretion from their nasus, which contains a mixture of terpenoids and elicits alarm and defensive recruitment (Thorne 1984; Roisin et al. 1990). Pheromones also communicate information about the presence and location of food, are used to recruit more soldiers and workers, and can modulate caste differentiation, either positively or negatively (Traniello and Busher 1985; Scharf 2015).

Recruitment was formerly proposed to depend on the quantitative production of the pheromone (Traniello and Busher 1985). In fact, it was effectively measured that the worker/soldier concentration ratio was 0.2 times for compounds concentrated in *N. corniger* (Arab et al. 2006). Moreover, behavioral evidence for species, colony, and caste specificity of the trail pheromone were also presented (Dunn and Messier 1999; Arab et al. 2006). In particular, Arab et al. (2006) described that both *N. corniger* workers and soldiers preferred the trails made by conspecific soldiers over those from workers. Extracts from the same species, and more particularly from the same colony, were also demonstrated to be preferred (Arab et al. 2006). Dunn and Messier (1999) highlighted that soldiers are able to discriminate their own colony members from conspecific neighbors and strangers; therefore, they behave more aggressively toward neighbors, with which they are competing for resources and territory, compared to strangers.

If quantitative differences between soldiers' and workers' trail pheromones can help to regulate the foraging behavior of these termites, the chemical composition of the sternal gland secretions should also be taken into account. For example, chemical analysis identified neocembrene, dodecatrienol and trinervitatriene in the pheromone trail, with a ratio of dodecatrienol/neocembrene of 1:25 and a ratio of trinervitatriene/neocembrene of 1:2 in the sternal glands of *N. corniger*'s workers (Sillam-Dussès et al. 2010). Monocyclic 14-membered ring diterpene cembrene-A is also a common constituent of this trail and is a short-term alarm capable of orienting soldiers to an area of disturbance (Hall and Traniello 1985). Moreover, some other parameters may play an important role in trail following. For example, a study performed by Gazal et al. highlighted the significant effect of feces on the trail (Gazal et al. 2014). Chemical analyses on the feces' composition, and behavioral studies implicating the identified compounds, may lead to valuable information that could help to improve the understanding of trail following mechanisms. This could be of importance to develop attractant or repulsive products in the context of an integrated pest management procedure. In addition to their crucial role for recruitment, chemical trails also probably exhibit antimicrobial activities and might be involved in the control of nest microbial competitors (Rosengaus et al. 2000). According to these results, studying the influence of the morphological type on potential specificity induction in trail preference could also lead to interesting results, besides contributing to the understanding of the social function of each morphological type inside a colony. Eventually, as primer pheromones from soldiers and other castes may also regulate caste homeostasis (Scharf et al., 2015), studying these compounds and their associated biology could potentially lead to insights for new control technologies.

Food selection: a complex process

Nasutitermes corniger feeds on dry, wet, or partially decomposed wood (He et al. 2013). It has the ability to consume a large variety of wood, such as dead twigs and branches in trees, tree stumps, wood debris on the ground and structural wood, including framing, plywood, pressboard, hardwood handles, wooden artwork, fences and paper and derivatives in laboratory conditions (Scheffrahn et al. 2005a). However, the exact stimuli implicated in the finding and selection of food by *N. corniger* is still under evaluation.

Nasutitermes corniger was first described as having no significant consumption preference among different wood species, but with a strong preference for decayed over fresh wood (Bustamante and Martius 1998). However, further contradictory studies demonstrated a significant species preference under laboratory conditions (Gazal et al. 2010). Yet, consistent with the results obtained by Bustamante and Martius (1998), the choice was not shown to exclusively depend on the density. Concerning the stage of decay of the wood, a more recent study conducted by Gazal et al. (2012) confirmed the results of Bustamante and Martius. Whatever the tested species, decayed wood always recruited more individuals than undecayed wood in standard foraging arenas. The next step was to investigate whether the choice of the source of wood was attributed to physical or chemical factors. This work thus demonstrated the crucial role of the secondary metabolites produced during wood degradation in the choice of a food substrate by comparing impregnation of a neutral (filter paper) and of wood (fresh or decayed) substrate with weathered or unweathered *Eucalyptus grandis* wood extracts (Gazal et al. 2014).

Following these results, further studies should be performed on the identification and origin of the compounds linked to wood degradation and to investigate their action on wood's physicochemical properties. It may also be interesting to investigate the effect of weathered and unweathered extracts on a neutral wood substrate from which secondary metabolites would have been removed or on non-durable wood, such as *Pinus* sp. According to some

authors, the preference for decayed wood could be due to several factors. Decomposition may lead to an increase of nutrient availability and a facilitation of mastication during the process and the depolymerization of cellulose, making wood fibers easier to remove and compounds more degradable. Moreover, it is possible that wood degradation releases allelochemicals that influence the foraging behavior or trigger trail following. Decomposition is also associated with the presence of microorganisms, notably fungi, which are potential additional sources of nitrogen and vitamins for the termites (Vasconcellos and Moura 2010; Gazal et al. 2012; Gazal et al. 2014). In addition to improve the ecological and biological knowledge of the species, understanding the mechanisms of termites' food selection would play a central role in developing new techniques to control *N. corniger* in case of infestation, such as substrates for insecticide-impregnated baits, or prevent *N. corniger* colonization of buildings by orienting technological choices.

Symbiosis and mutualism

The insect gut is divided into three regions: the stomodeum (foregut), the mesenteron (midgut), and the proctodeum (hindgut). The foregut includes the crop, a food-storage organ. In termites, the hindgut is enlarged and subdivided into five segments (P1-P5). The crop is slightly acidic, whereas the hindgut is moderately alkaline, with anaerobic conditions (Bignell and Anderson 1980). In the third hindgut segment (P3 or paunch), a bacterial community has developed that confers certain important benefits to the termite. In fact, in the case of *N. corniger*, such as other higher termites (Termitidae), digestive enzymes are encoded both by the termite genome itself and by these symbiotic bacteria, as these species do not possess hindgut flagellate protozoa, contrary to lower termites for which these organisms are the sources of cellulases and hemicellulases (Warnecke et al. 2007). However, this microbiome appears to play a far more complex role than only being related to cellulose degradation, as in

nitrogen fixation (Prestwich and Bentley 1981; Tayasu et al. 1994). In the case of an arboreal species close to *N. corniger*, a metagenomic analysis of the hindgut-associated microbial community highlighted the presence of bacterial genes for cellulose and xylan hydrolysis and also provided information about the possible functional role of this community concerning free H₂ generation and CO₂-reductive acetogenesis, alongside the identification of genes associated with nitrogen fixation (Warnecke et al. 2007). Metaproteome analysis of this community revealed the presence of proteins of known enzymatic functions in the case of *N. corniger*, thus permitting the reconstruction of the enzymatic pathways concerning transport, glucan binding, sugar fermentation, hydrogen metabolism, amino acid synthesis, and nitrogen fixation and assimilation (Burnum et al. 2010). These findings are complementary to the results previously obtained by Warnecke et al. (2007), although from all the isolated known proteins no enzyme involved in cellulose degradation could be observed. This may be due to methodic limitations, or because the cellulose native degradation may occur elsewhere than inside the termite hindgut.

Spirochetes, mostly from the *Treponema* genus, and Fibrobacteres, alongside an undetermined phylum (TG3), were the most abundant phyla of these *N. corniger* microbial symbionts and were established to be fiber-associated bacterial communities that significantly contribute to the cellulotic activity (He et al. 2013; Warnecke et al. 2007; Mikaelyan et al. 2014). Warnecke et al. (2007) also demonstrated that gene modules coding for iron-only hydrogenase were bound to the genus *Treponema*, as well as marker genes linked to the acetogenesis pathway, and this process apparently dominates in *Nasutitermes* by Spirochaetes. In the actinobacterial community, well known for its role in organic matter decomposition, such as cellulose, five families (*Propionibacteriaceae*, *Streptomycetaceae*, *Cellulomonadaceae*, *Corynebacteriaceae* and *Rubrobacteraceae*) were identified for *N. corniger*, regardless of the geographical origin or feeding behavior (Lefebvre et al. 2009).

Regarding the apparent role of microbial symbionts concerning cellulose digestion and other essential metabolic functions, disrupting this effective equilibrium would be important in the continuous search for insecticidal products. In fact, it was demonstrated that modifying the carbon source allocated to the higher termite *N. takasagoensis* led to marked changes in the bacterial community structure, thus highlighting the delicateness of this relationship (Miyata et al. 2007). Moreover, in the case of an integrated pest management strategy, searching for products that exhibit antimicrobial activity seems to be a promising perspective (Shelton and Grace 2003).

Concurrent to this crucial symbiosis, other types of associations can be observed in termites' societies. In particular, ants could be associated with facultative or opportunistic mutualism where these insects use part of the termite nest, prey on live or dead workers, provide the termites with nitrogen nutrients and assist in nest protection. The frequency of *N. corniger* association with ant species depends, in particular, on the cost-benefit balance linked to predation risk. A Brazilian study demonstrated that Formicidae species richness in *Nasutitermes*' nests (*N. corniger* and *N. ephratae*) was significantly higher in abandoned nests, in 48 species, compared to active or decadent nests, in 6 and 8 species (Santos et al. 2010). In the latter case, *Crematogaster acuta* (Fabricius) and *Ectatomma tuberculatum* (Olivier) were the most frequent species observed. *Monacis bispinosa* (Olivier), which is highly susceptible to the soldier frontal gland mixture and which do not prey on live termites, was the most frequently found species in common ants / *N. corniger* nests in Venezuela (Jaffe et al. 1995). *Crematogaster brevispinosa* (Mayr), an arboreal ant, has been observed in association with *N. corniger* in Brazil, sharing the same nest even if a physical barrier is built between the two communities. The participation of ants in nest protection and nutritional substrate supply thus appears to be another factor contributing to the success of *N. corniger* colonization.

437

438 **3.2 Pest management**

439 Synthetic chemical products, as active substances inhibiting wood degradation by termites
440 and microorganisms, can be applied through soil, wood treatment, or chemical fumigations,
441 before or after infestation (Paiva et al. 2011) (Table 1). Excellent reviews or technical reports,
442 such as those by Schultz et al. (2007; 2008) concerning wood preservation, Verma et al.
443 (2009) on the biological control of termites, or the expert group on termite biology and
444 management of the United Nations Environment Program (UNEP 2000), have taken an
445 inventory of the different strategies, depending on the geographical zone. Despite the
446 perceived efficiency of all of these substances, we cannot ignore the constraints on use,
447 environmental risk, human health and toxicity to non-target organisms (Pluke et al. 1999;
448 Clausen and Yang 2007; Santana et al. 2010; Scheffrahn and Scherer 2013; Scheffrahn et al.
449 2014). However, the efficacy of the different strategies mentioned above is mainly evaluated
450 against subterranean termite species. Despite their recognized status as a significant urban
451 pest, few references specifically mention arboreal termites, particularly *Nasutitermes* spp.
452 This observation can be explained by several factors. First, damages caused by subterranean
453 termites are less predictable, and preventive treatments must therefore be very efficient and
454 are crucially needed. In fact, in regard to physical management, the easily detected *N.*
455 *corniger* nests are often simply collected, broken and/or burnt (UNEP 2000; Pluke et al.
456 1999), whereas barriers often containing insecticides must be used as a management strategy
457 for subterranean termites (Verma et al. 2009; UNEP 2000; Paiva et al. 2011). A second factor
458 to be taken into account is the fact that *Nasutitermes* species mostly impact Central and South
459 America, alongside other tropical zones of the globe, such as Southeast Asia and Australia,
460 while the main commercial markets related to insecticidal products are located in the
461 Northern hemisphere, mostly concerned with subterranean termites (Constantino et al. 2002;

UNEP 2000; Miura et al. 2000). For example, in 2008, North America represented 60% of the total market for wood preserving biocides and Europe and Asia/Oceania represented 20% each (Schultz et al. 2008).

However, products for wood protection commercially available in the market must be active against a wide range of organisms, from decaying fungi to various xylophagous insects (Schultz et al. 2007). Even if species specificities exist, the insecticidal products mainly tested on subterranean termites (which target the workers caste) are thus also efficient towards arboreal termites, such as *N. corniger*. In the following sections, we thus focused as much as possible on environmentally low-risk management strategies, presenting examples specifically concerning *N. corniger* but also some references concerning *Nasutitermes* sp. and sometimes about other termites species when needed.

Botanical insecticides: antitermitic extracts and compounds

Numerous laboratory studies target the evaluation of botanical extracts and natural compounds against termites. Even if they mostly address the activity of these products against subterranean species, some interesting results can be cited concerning *Nasutitermes* spp. and, more particularly *N. corniger*.

One strategy for finding new specific termiticidal compounds from botanical sources is called “bioinspiration,” which aims to understand the functional role of secondary metabolites in living organisms, thus gaining inspiration from the abilities of plants or microorganisms to produce adapted bioactive molecules under specific environmental conditions (Rodrigues et al. 2010; Basset et al. 2012; Houël et al. 2015). In particular, one source for these antitermitic natural compounds are woods naturally resistant to biodegradation, which is closely linked to the accumulation of extractives typically in the heartwood (Taylor et al. 2002; Ragon et al. 2008; Kampe and Magel 2013). Results

concerning wood extracts activity against *N. corniger* or *Nasutitermes* sp. (Peres Filho et al. 2006; Barbosa et al. 2007; Sá et al. 2008; 2009; Santana et al. 2010; Boulogne et al. 2011; Rodrigues et al. 2011; Rodrigues et al. 2010) are summed up in Table 2.

The research for natural extracts for wood protection contemplates various modes of action such as elicitors, regulators, inhibitors, enzymes, or repellents for pest control. In fact, specific defense mechanisms were evidenced in plant seeds and roots, which are essential for the propagation and survival of the species. Lectins, for example, which are toxic proteins widely distributed among plants, are supposedly involved in defense mechanisms against phytophagous predators (Carlini and Grossi 2002). These compounds, isolated from various sources, were extensively studied for their insecticidal activity, *N. corniger* being among the selected targets (Sá et al. 2008; Silva et al. 2009; Napoleão et al. 2011; Paiva et al. 2011; Souza et al. 2011). These proteins play a role in cell agglutination by carbohydrate recognition and could therefore act by binding to glycoproteins present in the insect's digestive tract (Sá et al. 2008). They were demonstrated to be able to disrupt enzymes' activities from the gut of termites (Lima et al. 2014). These molecules are now the basis of commercial formulations currently developed against termites (Paiva et al. 2011). Eventually, some well-known insecticidal compounds from plant extracts are active as biocides, strong repellants or antifeedants against several species, including subterranean termites (Sharma et al. 1991; Grace and Yates 1992; Serit et al. 1992; Ivbijaro 1993; Delate and Grace 1995; Chuanchuan et al. 1998; UNEP 2000; YunLing et al. 2009; Tahiri et al. 2011; Sharma et al. 2011; Himmi et al. 2013; Boulogne et al. In prep.) (Table 2).

Concurrently to the evaluation of extracts' biological activity, several reviews have been reported in natural products for wood protection (Singh and Singh 2012; Verma et al. 2009; Yang 2009; Mai et al. 2004; Kirker et al. 2013; González-Laredo et al. 2015). These reports discuss the development of botanical pesticides against subterranean termites, with in

particular results concerning *Nasutitermes* wood decay control. Some extracts could be employed unadjuvanted as wood protectants, as reported in an African study, about the effective extracts of *Erythrophleum suaveolens* (Guill. & Perr.) Brenan (Fabaceae), *Azadirachta indica* A. Juss. (Meliaceae) and *Chromolaena odorata* (L.) King & Rob. (Asteraceae) in improving the durability of *Antiaris toxicaria* Lesch. (Moraceae) wood, exposed to decay for five years (Antwi-Boasiako and Damoah, 2010). In another study, *Cryptomeria japonica* (Thunb. ex L. f.) D. Don (Cupressaceae) wood treated with essential oils, showed strong activity against subterranean termites, in a three weeks test (Kartal *et al.*, 2006). Turner and Conradie have also demonstrated the enhanced termitic resistance of *Pinus patula* Schltdl. & Cham. (Pinaceae) treated with extracts from four African durable species - *Breonadia salicina* (Vahl) Hepper & Wood (Rubiaceae), *Spirostachys Africana* Sond. (Euphorbiaceae), *Ptaeroxylon obliquum* (Thunb.) Radlk. (Rutaceae) and *Syncarpia glomulifera* (Sm.) Nied. (Myrtaceae)- in a two years test (Turner and Conradie, 1995).

Another way to employ botanical pesticides is to combine natural products to other protectants compounds, such as described in a 2007 study, where 0.5 percent thujaplicin was used in several multicomponent biocide systems with borate and carbon-based biocides. Noticeable results were obtained for protecting wood from mold, decay, and termite damage (Clausen and Yang 2007).

Despite all these advances, until now there has been no available chemical formulation containing these compounds for wood treatment against termites. Moreover, although they present a repulsive effect against *N. corniger*, their effects are not persistent and they did not lead to any contact toxicity (Paes et al. 2010).

Botanical insecticides: antimicrobial extracts and compounds

In most previously cited studies, the authors chose to use bioassays using the whole insect as a target, thus highlighting potent extracts or molecules. However, another process could be used to precisely search for compounds or extracts acting on a selected mechanism of action, as exemplified by Bulmer et al. (2009) with the glycomimetic molecule D- δ -gluconolactone, as above cited in this review. Thus, in-depth studies of *N. corniger* biology could be helpful for the discovery of targeted alternatives to non-specific pesticides.

Another specific approach based on natural processes is searching for phytochemicals targeting termites bacterial gut community. Even if some publications describe antimicrobial activities of compounds isolated from *Nasutitermes* sp. (Zhao et al. 2004; Siderhurst et al. 2005; Coutinho et al. 2009; Coutinho et al. 2010; Nirma et al. 2013), few articles describe the search for antibacterial products according to the perspective of an integrated pest management strategy against *Nasutitermes* species. However, a recent study showed that botanical lectins had an antibacterial effect on symbiotic bacteria from the gut of *N. corniger* (Napoleão et al. 2011). This approach could be of valuable interest, as similar results have also been achieved with other termite species (Table 2) (Maistrello et al. 2001; Maistrello et al. 2003; Ahmed et al. 2006; Doolittle et al. 2007).

Other biopesticides

Other strategy for finding new termiticidal compounds from other sources is considering soil actinomycetes fermentation products. These naturally occurring compounds were discovered from *Streptomyces* sp. culture broths. Some studies dealt with control of termites with these pesticides. Indeed, Chen et al. (2015) found that avermectin (abamectin and ivermectin) has optimal insecticidal properties to manage the subterranean termite *Reticulitermes speratus*. This strategy could be tested in the control of *Nasutitermes* but it has to be taken into account that these compounds have a large range of action and thus

potentially noticeable impact in the field, being used in particular as an agent against worms, ticks, flies and other insects in veterinary applications (Duke et al. 2010).

Biological control of termites: entomopathogens and natural enemies

Nematodes, bacteria, and fungi have several interesting properties for the biological control of termites (Verma et al. 2009; Paiva et al. 2011). In recent years, the number of publications concerning the biological control of termites has increased dramatically (Chouvenc et al. 2011). In a critical review based on the study of 227 publications in which 427 bioassays were reported, Chouvenc et al. (2011) showed that these strategies were largely unsuccessful. Even if numerous examples of bacterial and nematode control can be found in the recent literature, mostly concerning subterranean termites (Grewal et al. 1999; Mo et al. 2006; Devi et al. 2007; Verma et al. 2009; Chouvenc et al. 2011), the authors raise questions about the fact that the described protocols may encounter poor biological relevancy, that the obtained data may often be improperly and overly optimistically interpreted, and that field bioassays often lack (Chouvenc et al. 2011). Other reasons that potentially explain this lacking of success may be linked to the remarkable way that termites have evolved to resist diseases due to biochemical and immunological strategies (Meikle et al. 2005), as also exemplified earlier in this review.

Among the few successes, biological control using fungi against *Nasutitermes* sp. (Table 2) should be mentioned (Ensaf et al. 2006; Chouvenc et al. 2011; Salas-Acuña 2012). A commercial formulation using the generalist entomopathogenic fungus *Metarhizium anisopliae* (formerly *flavoviride*) (Metsch.) Sorokin (BioblastTM, Ecoscience Co.) was developed, but was unfortunately removed from the market in the early 2000s due to insufficient results (Verma et al. 2009; Chouvenc et al. 2011). The pathogenicity of the fungus is dependent on various factors, such as its capacity for germination and penetration

on and through the cuticle as well as the insect's immune system to prevent fungus growth (Verma et al. 2009). A perspective to improve the efficacy of this enthomopathogen would therefore be to collect strains directly from native infected termites to improve the probability of isolating fungi having circumvented termites' adaptation strategies and natural defenses (Meikle et al. 2005). The challenge in termites' biological control is thus to develop more biologically relevant models, promote interdisciplinarity to finalize and conduct the assays, and add field and technology transfer studies (Chouvenc et al. 2011). As described earlier in their review, Chouvenc et al. (2011) also clearly appeal to an increase in research concerning termites' biology to find ways for pathogens to bypass termites' defense mechanisms.

In addition to entomopathogens, other biological control agents may involve some predators like ants, such as *Camponotus abdominalis*, one of the most aggressive species of this fierce natural enemy of termites (UNEP 2000). Another well-known predator of *N. corniger* is the hemipteran *Salyavata variegata* nymph (Reduviidae), which captures termite workers with the help of traps made of empty carcasses of previous prey and carton crumbs (McMahan 1982). However, the investigation of natural predators as control agents did not reveal any commercial potential (Chouvenc et al. 2011).

3.3 Not only a pest!

As with other termite species, *N. corniger* is a decomposer insect that plays a fundamental role in the decomposition process and nutrient cycling by aerating the soil, recycling plant matter and fixing nitrogen (Vasconcellos and Moura 2010). The presence of these insects also indicates the good health of an ecosystem, and they can be considered as bioindicators of its actual function (Täyasu et al. 1994; Paiva et al. 2011). Moreover, the presence of termites has a positive impact on biodiversity, termitariophile organisms, or insects depending on the nest structure for shelter, food or reproduction, thus being

maintained in the environment (Jaffe et al. 1995; Santos et al. 2010). Eventually, arboreal nests may also provide homes for other animals, such as birds and bats (Postava-Davignon et al. 2010).

Concurrently with playing a role in dynamic environmental processes, *N. corniger* can also provide services to mankind, acting as a source of active secondary metabolites, particularly antimicrobial compounds. Inspired from its traditional use as a medicine in Northeastern Brazil, aqueous extracts from *N. corniger* were studied and shown to exhibit low antimicrobial activity against *Escherichia coli* aminoglycoside-resistant strains, but presented interesting synergist activities with antibiotics (Coutinho et al. 2009; 2010). Extracts from *N. corniger* could therefore be used as sources of compounds to bypass bacterial resistance mechanisms. Other results highlighted antibacterial (Zhao et al. 2004) or antifungal (Bulmer and Crozier 2004) activities for compounds isolated from *Nasutitermes* species. Eventually, active compounds can also be obtained through termites' symbionts, such as the antifungal compounds tyroscherin and N-methyltyroscherin, isolated from *Pseudallescheria boydii*, a saprophytic fungus frequently found in soil and water, but that was isolated in this case from *Nasutitermes* sp. (Nirma et al. 2013). In addition to appealing to the preservation and the pharmacochemical study of this species, these examples highlight the fact that improving the knowledge of termites, and more particularly *N. corniger* biology, could lead to a marked improvement, not only in the pest management field but also in the continuous search for new antimicrobial compounds.

4. Conclusion

Nasutitermes is the most species-rich termite genus, with a complex taxonomic history, and a growing number of known species present in all biogeographical regions. *Nasutitermes corniger*, which is the type-species of the genus, is an urban and agricultural

pest, and also an invasive species, with a remarkable ability to adapt to a variety of habitats and food sources. Controversially, ecological and human health problems potentially arising from the use of synthetic chemical substances, particularly if the label guidelines are not strictly followed, are the limiting factor for their use. However, the extensive use of botanical or biological insecticides remains up to date inconclusive because of their ineffectiveness or their poor durability in external wood exposure cases. In the continuous effort to develop safe and effective methods for conehead termite management, we therefore believe that a paradigm shift is necessary and that sustained efforts must be performed in the field of termite biology. Alongside an increase in field bioassays, improvement in this knowledge is the key to more elaborate pertinent models and integrated strategies, targeting both the insect and its symbionts.

Eventually, integrated management rather than eradication is necessary in the native zones of occurrence of this termite. Although it is considered to be a pest, the termite is clearly an essential component of ecosystems as well as a potential source of valuable compounds that could contribute to the improvement of human health. Thus, the perspective remains open concerning *Nasutitermes* species study and management, and exciting discoveries are still to be made.

Acknowledgements

The authors gratefully thank Dr. Philippe Petit (French West Indies University, UA) for providing *Lonchocarpus* sp. vegetal material, and Arnaud Jahn-Oyac for his technical help. The authors also wish to thank Dr. Thomas Chouvenc from University of Florida for identifying *N. corniger* from our field collections. This work is part of the INSECTICIDES project funded by Europe (ERDF OP), French Guiana Regional Council and the Air Liquide

Foundation. This work benefited from an "Investissement d'Avenir" grant managed by Agence Nationale de la Recherche (CEBA, ref. ANR-10-LABX-25-01).

References

Adams E, Atkinson L (2008) Queen fecundity and reproductive skew in the termite *Nasutitermes corniger*. Insect Soc 55:28-36

Ahmed S, Riaz MA, Shahid M (2006) Response of *Microtermes obesi* (Isoptera: Termitidae) and its gut bacteria towards some plant extracts. J Food Agric Environ 4:317-320

Arab A, Issa S, Alfonzo D, Jaffe K (2006) Caste, colony, and species specificity of the trail pheromone in two sympatric Nasutitermitinae (Isoptera: Termitidae). Sociobiology 47:345-351

Atkinson L, Adams ES (1997) The origins and relatedness of multiple reproductives in colonies of the termite *Nasutitermes corniger*. P Roy Soc Lond B Bio 264:1131-1136

Atkinson L, Teschendorf G, Adams ES (2008) Lack of evidence for nepotism by workers tending queens of the polygynous termite *Nasutitermes corniger*. Behav Ecol Sociobiol 62:805-812

Banks N (1918) The termites of Panama and British Guiana. B Am Mus Nat Hist 38: 659–667

- Barbosa AP, Nascimento CS, Morais JW (2007) Studies on the antitermitic properties of wood and bark crude extracts of forest species from Central Amazonian, Brazil. *Acta Amazonica* 37:213-218
- Basset C, Rodrigues AMS, Eparvier V, Silva MRR, Lopes NP, Sabatier D, Fonty E, Espindola LS, Stien D (2012) Secondary metabolites from *Spirotropis longifolia* (DC) Baill and their antifungal activity against human pathogenic fungi. *Phytochemistry* 74: 166-172.
- Bignell D, Anderson J (1980) Determination of pH and oxygen status in the guts of lower and higher termites. *J Insect Physiol* 26:183-188
- Boulogne I, Germosen-Robineau L, Ozier-Lafontaine H, Jacoby-Koaly C, Aurela L, Loranger-Merciris G (2011) *Acromyrmex octospinosus* (Hymenoptera: Formicidae) management. Part1. Effects of TRAMILs insecticidal plant extracts. *Pest Manage Sci* 68:313-320
- Boulogne I, Falkowski M, Jahn Oyac A, Dusfour I, Houël E, Petit P (*in prep*) Investigating a strong difference of insecticidal activity and cytotoxicity between two thinly known ichthyotoxic species: *Lonchocarpus chrysophyllus* Kleinhoonte and *Lonchocarpus floribundus* Benth
- Brauer F (1868) Verzeichniss der bis jetzt bekannten Neuropteren im Sinne Linné's. *Verhandlungen der ZoologischBotanischen Gesellschaft in Wien* 18: 359–416

- Brune A (2014) Symbiotic digestion of lignocellulose in termite guts. *Nat Rev Microbiol* 12:168-180
- Bulmer M S, Crozier RH (2004) Duplication and diversifying selection among termite antifungal peptides. *Mol Biol Evol* 21:2256-2264
- Bulmer MS, Bachelet I, Raman R, Rosengaus RB, Sasisekharan R (2009) Targeting an antimicrobial effector function in insect immunity as a pest control strategy. *P Roy Soc Lond B Bio* 106:12652-12657
- Burnum KE, Callister SJ, Nicora CD, Purvine SO, Hugenholtz P, Warnecke F, Scheffrahn RH, Smith RD, Lipton MS (2010) Proteome insights into the symbiotic relationship between a captive colony of *Nasutitermes corniger* and its hindgut microbiome. *ISME J* 5:161-164
- Bustamante NC and Martius C (1998) Nutritional attractiveness of wood-feeding termites inhabiting floodplain forest of the Amazon river, Brazil. *Acta Amazonica* 28:301-307
- Carlini C, Grossi-de-Sa MF (2002) Plant toxic proteins with insecticidal properties. A review on their potentialities as bioinsecticides. *Toxicon* 40:1515-1539
- Chen Z, Qu Y, Xiao D, Song L, Zhang S, Gao X, Song D (2015) Lethal and social-mediated effects of ten insecticides on the subterranean termite *Reticulitermes speratus*. *J Pest Sci* 88:741-751

- Chouvenc T, Su N-Y, Kenneth Grace J (2011) Fifty years of attempted biological control of termites - Analysis of a failure. Biol Control 59:69-82
- Chuanchuan L, Rongjing X, Haoquan Z, Keng C (1998) The effects of a Neem insecticide on orientation and feeding behaviour of *Coptotermes formosanus* (Isoptera:Rhinotermitidae).http://en.cnki.com.cn/Article_en/CJFDTOTALBYKJ199801001.htm
- Clausen CA, Yang V (2007) Protecting wood from mould, decay, and termites with multi-component biocide systems. Int Biodeter Biodegr 59:20-24
- Constantino R (2002) The pest termites of South America: taxonomy, distribution and status. J Appl Entomol 126:355-365
- Constantino R (2000) Online termite database. Departamento de Zoologia, Universidade de Brasília. <http://www.unb.br/ib/zoo/docente/constant/catal/catnew.html>
- Coutinho HD, Vasconcellos A, Freire-Pessôa HL, Gadelha CA, Gadelha TS, Almeida-Filho GG (2010) Natural products from the termite *Nasutitermes corniger* lowers aminoglycoside minimum inhibitory concentrations. Pharmacogn Mag 6:1-4
- Coutinho HD, Vasconcellos A, Lima MA, Almeida-Filho GG, Alves RR (2009) Termite usage associated with antibiotic therapy: enhancement of aminoglycoside

antibiotic activity by natural products of *Nasutitermes corniger* (Motschulsky 1855).
BMC Complem Altern M 9:35

Cuezzo C, Carrijo TF, Canello EM (2015) Transfer of two species from *Nasutitermes*
Dudley to *Cortaritermes* Mathews (Isoptera: Termitidae: Nasutitermitinae). Austral
Entomol 54: 172–179

Delate KM, Grace JK (1995) Susceptibility of neem to attack by the Formosan
subterranean termite, *Coptotermes formosanus* Shir. (Isopt., Rhinotermitidae). J Appl
Entomol 119:93-95

Devi KK, Seth N, Kothamasi S, Kothamasi D (2007) Hydrogen cyanide-producing
rhizobacteria kill subterranean termite *Odontotermes obesus* (rambur) by cyanide
poisoning under in vitro conditions. Curr Microbiol 54:74-78

Doolittle M, Raina A, Lax A, Boopathy R (2007) Effect of natural products on gut
microbes in Formosan subterranean termite, *Coptotermes formosanus*. Int Biodeter
Biodegr 59:69-71

Dunn R, Messier S (1999) Evidence for the opposite of the dear enemy phenomenon
in termites. J Insect Behav 12:461-464

Dudley PH (1890) Termites of the Isthmus of Panama. Part II. Transactions of the New
York Academy of Science 9:157–180

- Duke SO, Cantrell CL, Meepagala KM, Wedge DE, Tabanca N, Schrader KK (2010)
Natural toxins for use in pest management. *Toxins* 2:1943-1962
- Eggleton P (2001) Termites and trees: a review of recent advances in termite
phylogenetics. *Insect Soc* 48:187-193
- Ensaf A, Bouree P, Callard P (2006) Potentialités de *Termitaria coronata* thaxter, 1920
(Deutéromycètes, Termitariales) pour la lutte biologique contre les termites
(Isoptera). *Bulletin de la Société entomologique de France* 111:43-49
- Evans TA, Forschler BT, Grace JK (2013) Biology of invasive termites: a worldwide
review. *Ann Rev Entomol* 58:455-474
- Fontes LR (1998) Novos aditamentos ao “Catálogo dos Isoptera do Novo Mundo”, e uma
filogenia para os gêneros neotropicais de Nasutitermitinae. In: L.R. Fontes & E. Berti-
Filho, Eds. *Cupins: O desafio do conhecimento*. FEALQ, Piracicaba, pp. 309–412
- Fontes LR, Milano S (2002) Termites as an urban problem in South America.
Sociobiology 40:103-151
- Froggatt WW (1897) Australian termites. Part II. *P Linn Soc N S W* 21:510–552
- Fuller CA, Postava-Davignon M (2014) Termites like it hot and humid: the ability of
arboreal tropical termites to mediate their nest environment against ambient
conditions. *Ecol Entomol* 39:253-262

- Fuller CA, Postava-Davignon M (2014) Termites like it hot and humid: the ability of arboreal tropical termites to mediate their nest environment against ambient conditions. *Ecol Entomol* 39:253-262
- Ganapaty S, Thomas PS, Fotso S, Laatsch H (2004) Antitermitic quinones from *Diospyros sylvatica*. *Phytochemistry* 65:1265-1271
- Gazal V, Bailez O, Viana-Bailez AM (2014) Mechanism of trail following by the arboreal termite *Nasutitermes corniger* (Isoptera: Termitidae). *Zool Sci* 31:1-5
- Gazal V, Bailez O, Viana-Bailez AM (2010) Wood preference of *Nasutitermes corniger* (Isoptera: Termitidae). *Sociobiology* 55:433-443
- Gazal V, Bailez O, Viana-Bailez AM, Aguiar-Menezes EL, Menezes EB (2012) Decayed wood affecting the attraction of the pest arboreum termite *Nasutitermes corniger* (Isoptera: Termitidae) to resource foods. *Sociobiology* 59:287-295
- Gazal V, Bailez O, Viana-Bailez AM, Aguiar-Menezes EL, Menezes EB (2014) Behavioral responses of the arboreal termite *Nasutitermes corniger* (Isoptera: Termitidae) to wood extracts. *Wood Sci Technol* 48:581-590
- Grace JK, Yates JR (1992) Behavioural effects of a neem insecticide on *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Tropical Pest Management* 38:176-180

834 Grewal P, Georgis R (1999) Entomopathogenic nematodes. In: Biopesticides: Use and
835 Delivery. Springer, pp. 271-299
836

837 Gush TJ, Bentley BL, Prestwich GD, Thorne BL (1985) Chemical variation in defensive
838 secretions of four species of *Nasutitermes*. Biochem Syst Ecol 13:329-336
839

840 Hagen HA (1858) Specielle Monographie der Termiten. Linnea Entomologica 12:4-342
841

842 Hall P, Traniello JF (1985) Behavioral bioassays of termite trail pheromones. J Chem
843 Ecol 11:1503-1513
844

845 Hartke TR, Rosengaus RB (2013) Costs of pleometrosis in a polygamous termite. P Roy
846 Soc Lond B Bio 280:1756
847

848 Harun J, Labosky Jr P (1985) Antitermitic and antifungal properties of selected bark
849 extractives. Wood Fiber Sci 17:327-335
850

851 He S, Ivanova N, Kirton E, Allgaier M, Bergin C, Scheffrahn RH, Kyrpides NC,
852 Warnecke F, Tringe SG, Hugenholtz P (2013) Comparative metagenomic and
853 metatranscriptomic analysis of hindgut paunch microbiota in wood-and dung-feeding
854 higher termites. PloS one 8:e61126
855

856 Himmi SK, Tarmadi D, Ismayati M, Yusuf S (2013) Bioefficacy performance of
857 neem-based formulation on wood protection and soil barrier against subterranean

858 termite, *Coptotermes gestroi* Wasmann (Isoptera: Rhinotermitidae). *Procedia*
859 *Environmental Sciences* 17:135-141

860

861 Holmgren N (1909) Termitenstudien 1. Anatomische untersuchungen. Kungliga Svenska
862 Vetenskapsakademiens Handlingar 44:1–215

863

864 Holmgren N (1910) Versuch einer Monographie der amerikanischen Eutermes - Arten.
865 Jahrbuch der Hamburgischen Wissenschaftlichen Anstalten 27:171–325

866

867 Holmgren N (1912) Termitenstudien 3. Systematik der Termiten. Die Familie
868 Metatermitidae. Kungliga Svenska Vetenskapsakademiens Handlingar 48:1–166

869

870 Houël E, Gonzalez, G, Bessière J-M, Odonne G, Eparvier V, Deharo E, Stien D (2015)
871 Therapeutic switching: from antidermatophytic essential oils to new leishmanicidal
872 products. *Mem I Oswaldo Cruz* 110:106-113

873

874 Inward DJG, Vogler AP, Eggleton P (2007) A comprehensive phylogenetic analysis
875 of termites (Isoptera) illuminates key aspects of their evolutionary biology. *Mol*
876 *Phylogenet Evol* 44: 953–967

877

878 ITIS (2014) from the Integrated Taxonomic Information System on-line database,
879 <http://www.itis.gov>. [Accessed May, 7, 2014]

880

881 Ivbijaro MF, Umeh V, Mutsaers H (1993) Laboratory toxicity of the crude extracts of
882 *Piper guineense* Schum and Thonn, *Azadirachta indica* A. Juss and *Parkia*

clappertoniana (Jacq) to the termites *Macrotermes nigeriensis* (Sjostedt) (Isoptera: Termitidae). Int J Trop Insect Sc 14:229-233

Jaffe K, Ramos C, Issa S (1995) Trophic interactions between ants and termites that share common nests. Ann Entomol Soc Am 88:328-333

Jones R (1979) Expansion of the nest of *Nasutitermes costalis*. Insectes Sociaux 26:322-342

Jones R (1980) Gallery construction by *Nasutitermes costalis*: Polyethism and the behavior of individuals. Insect Soc 27:5-28

Kampe A, Magel E (2013) New insights into heartwood and heartwood formation. In: Cellular Aspects of Wood Formation. Springer Berlin Heidelberg, pp 71-95

Krishna K, Grimaldi DA, Krishna V, Engel MS (2013) Treatise on the Isoptera of the World. B Am Mus Nat Hist 377: 1-2704

Lefebvre T, Miambi E, Pando A, Diouf M, Rouland-Lefèvre C (2009) Gut-specific actinobacterial community structure and diversity associated with the wood-feeding termite species, *Nasutitermes corniger* (Motschulsky) described by nested PCR-DGGE analysis. Insect Soc 56:269-276

Levings SC, Adams ES (1984) Intra-and interspecific territoriality in *Nasutitermes* (Isoptera:Termitidae) in a Panamanian mangrove forest. J Anim Ecol 53:705-714

- Lima VLS, Bailez OE, Viana-Bailez AM (2013) Caste polymorphism of apterous line of the Neotropical termite *Nasutitermes corniger* (Motschulsky) (Isoptera, Termitidae). Rev Bras Entomol 57:309-312
- Lima T de A, Pontual EV, Dornelles LP, Amorim PK, Sá RA, Coelho LCBB, Napoleão TH, Paiva PMG (2014) Digestive enzymes from workers and soldiers of termite *Nasutitermes corniger*. Comp Biochem Physiol Part B: Biochem Mol Biol 176:1-8
- Linnaeus C (1758) Systema Naturae. 10th ed. Laurentii Salvii, Holmiae
- Maistrello L, Henderson G, Laine RA (2003) Comparative effects of vetiver oil, nootkatone and disodium octaborate tetrahydrate on *Coptotermes formosanus* and its symbiotic fauna. Pest Manage Sci 59:58-68
- Maistrello L, Henderson G, Laine RA (2001) Efficacy of vetiver oil and nootkatone as soil barriers against Formosan subterranean termite (Isoptera: Rhinotermitidae). J Econ Entomol 94:1532-1537
- Mathews AGA (1977) Studies on Termites from the Mato Grosso State, Brazil. Academia Brasileira de Ciências, Rio de Janeiro
- McMahan E (1982) Bait-and-capture strategy of a termite-eating assassin bug. Insect Soc 29:346-351

- McMahan EA (1970) Polyethism in workers of *Nasutitermes costalis* (Holmgren).
Insect Soc 17:113-120
- Meikle WG, Mercadier G, Rosengaus RB, Kirk AA, Derouané F, Quimby PC (2005)
Evaluation of an entomopathogenic fungus, *Paecilomyces fumosoroseus* (Wize)
Brown and Smith (Deuteromycota: Hyphomycetes) obtained from Formosan
subterranean termites (Isop., Rhinotermitidae). J Appl Entomol 129:315-322
- Mello AP, Costa BG, Santos AC, Silva AMB, Bezerra-Gusmão MA (2014) Termite
infestation in historical buildings and residences in the semiarid region of Brazil.
Sociobiology 61:318-323
- Mikaelyan A, Strassert JFH, Tokuda G, Brune A (2014) The fibre-associated cellulolytic
bacterial community in the hindgut of wood-feeding higher termites (*Nasutitermes*
spp.). Environ Microbiol 16:2711-2722
- Miura T, Roisin Y, Matsumoto T (2000) Molecular phylogeny and biogeography of the
Nasute termite genus *Nasutitermes* (Isoptera: Termitidae) in the Pacific Tropics. Mol
Phylogenet Evol 17:1-10
- Miyata R, Noda N, Tamaki H, Kinjyo K, Aoyagi H, Uchiyama H, Tanaka H (2007)
Influence of feed components on symbiotic bacterial community structure in the gut of
the wood-feeding higher termite *Nasutitermes takasagoensis*. Biosci Biotech Bioch
71:1244-1251

- Mo J, Wang Z, Song X, Guo J, Cao X, Cheng J (2006) Effects of sublethal concentration of ivermectin on behaviour of *Coptotermes formasanus*. *Sociobiology* 47:687-696
- Müller F (1873) Beiträge zur Kenntniss der Termiten I-II. *Jenaische Zeitschrift für Medizin und Naturwissenschaft* 7:333–358
- Napoleão TH, Gomes FS, Lima TA, Santos ND, Sá RA, Albuquerque AC, Coelho LC, Paiva PM (2011) Termiticidal activity of lectins from *Myracrodruon urundeuva* against *Nasutitermes corniger* and its mechanisms. *Int Biodeter Biodegr* 65:52-59
- Nirma C, Eparvier V, Stien D (2013) Antifungal agents from *Pseudallescheria boydii* SNB-CN73 isolated from a *Nasutitermes* sp. termite. *J Nat Prod* 76:988-991
- Nirma C, Rodrigues AM, Basset C, Chevolot L, Girod R, Moretti C, Stien D, Dusfour I, Eparvier V (2012) Larvicidal activity of isoflavonoids from *Muelleria frutescens* extracts against *Aedes aegypti*. *Nat Prod Comm* 7:1319-1322
- Paes JB, de Souza AD, de Lima CR, de Medeiros Neto PN (2010) Efficiency of neem and castor oil plant oils against xylophagous termites under forced food assay. *Cerne* 16:105-113
- Paiva PM, Santana G, Souza IF, Albuquerque LP, Agra-Neto AC, Albuquerque AC, Luz LA, Napoleão TH, Coelho LC (2011) Effect of lectins from *Opuntia ficus indica* cladodes and *Moringa oleifera* seeds on survival of *Nasutitermes corniger*. *Int Biodeter Biodegr* 65:982-989

983

984 Peres Filho O, Dorval A, Duda MJ, Moura RG (2006) *Nasutitermes* sp. (Isoptera,

985 Termitidae) response to extract from four brazilian woods. *Scientia Florestalis* 71:51-

986 54

987

988 Pluke RWH, Permaul D, Leibee GL (1999) Integrated pest management and the use of

989 botanicals in Guyana. Bib. Orton IICA/CATIE

990

991 Postava-Davignon MA (2010) Evolution and ecology of termite nesting behavior and its

992 impact on disease susceptibility. Biology Dissertations

993 <http://hdl.handle.net/2047/d20000126>

994

995 Prestwich G and Bentley BL (1981) Nitrogen fixation by intact colonies of the termite

996 *Nasutitermes corniger*. *Oecologia* 49:249-251

997

998 Prestwich GD (1979) Chemical defense by termite soldiers. *J Chem Ecol* 5:459-480

999

1000 Ragon KW, Nicholas DD, Schultz TP (2008) Termite-resistant heartwood: The effect

1001 of the non-biocidal antioxidant properties of the extractives (Isoptera:

1002 Rhinotermitidae). *Sociobiology* 52:47-54

1003

1004 Rodrigues A (2010) Valorisation des extractibles responsables de la durabilité naturelle

1005 des bois exploités de Guyane Française. PhD Thesis, Université des Antilles et de la

1006 Guyane and Université fédérale de Brasilia

1007

1008 Rodrigues A, Amusant N, Beauchêne J, Eparvier V, Leménager N, Baudassé C,
 1009 Espindola LS, Stien D (2011) The termiticidal activity of *Sextonia rubra* (Mez) van
 1010 der Werff (Lauraceae) extract and its active constituent rubrynolide. Pest Manage Sci
 1011 67:1420-1423
 1012
 1013 Rodrigues AMS, Theodoro PNET, Eparvier V, Basset C, Silva MRR, Beauchêne J,
 1014 Espindola LS, Stien D (2010) Search for antifungal compounds from the wood of
 1015 durable tropical tress. J Nat Prod 73:1706-1707
 1016
 1017 Roisin Y, Everaerts C, Pasteels J, Bonnard O (1990) Caste-dependent reactions to soldier
 1018 defensive secretion and chiral alarm/recruitment pheromone in *Nasutitermes princeps*.
 1019 J Chem Ecol 16:2865-2875
 1020
 1021 Rojas MG and Morales-Ramos JA (2001) Bait matrix delivery of chitin synthesis
 1022 inhibitors to the Formosan subterranean termite (Isoptera: Rhinotermitidae). J Econ
 1023 Entomol 2:506-510
 1024
 1025 Rosengaus RB, Lefebvre ML, Traniello JF (2000) Inhibition of fungal spore germination
 1026 by *Nasutitermes*: evidence for a possible antiseptic role of soldier defensive secretions.
 1027 J Chem Ecol 26:21-39
 1028
 1029 Rouland-Lefèvre C (2011) Termites as Pests of Agriculture. In: Bignell ED, Roisin Y, Lo
 1030 N Eds, Biology of Termites: A Modern Synthesis. Springer Netherlands, Dordrecht,
 1031 pp. 499–517
 1032

1033 Roy V, Constantino R, Chassany V, Giusti Miller S, Diouf M, Mora P, Harry M (2014)
 1034 Species delimitation and phylogeny in the genus *Nasutitermes* (Termitidae:
 1035 *Nasutitermitinae*) in French Guiana. *Mol Ecol* 23:902-920
 1036
 1037 Sá RA, Argolo AC, Napoleão TH, Gomes FS, Santos ND, Melo CM, Albuquerque AC,
 1038 Xavier HS, Coelho LC, Bieber LW (2009) Antioxidant, *Fusarium* growth
 1039 inhibition and *Nasutitermes corniger* repellent activities of secondary metabolites
 1040 from *Myracrodruon urundeuva* heartwood. *Int Biodeter Biodegr* 63:470-477
 1041
 1042 Sá RA, Napoleão TH, Santos ND, Gomes FS, Albuquerque AC, Xavier HS, Coelho LC,
 1043 Bieber LW, Paiva PM (2008) Induction of mortality on *Nasutitermes corniger*
 1044 (Isoptera, Termitidae) by *Myracrodruon urundeuva* heartwood lectin. *Int Biodeter*
 1045 *Biodegr* 62:460-464
 1046
 1047 Salas-Acuña B (2012) Posibilidades de control de *Nasutitermes corniger* utilizando el
 1048 hongo entomopatógeno *Metarhizium* sp. *Revista Forestal Mesoamericana Kurú* 2:16
 1049
 1050 Santana AL, Maranhão CA, Santos JC, Cunha FM, Conceição GM, Bieber LW,
 1051 Nascimento MS (2010) Antitermitic activity of extractives from three Brazilian
 1052 hardwoods against *Nasutitermes corniger*. *Int Biodeter Biodegr* 64:7-12
 1053
 1054 Santos PP, Vasconcellos A, Jahyny B, Delabie JHC (2010) Ant fauna (Hymenoptera,
 1055 Formicidae) associated to arboreal nests of *Nasutitermes* spp. (Isoptera, Termitidae) in
 1056 a cacao plantation in southeastern Bahia, Brazil. *Rev Bras Entomol* 54:450-454
 1057

1058 Scharf ME (2015) Termites as targets and models for biotechnology. *Annu Rev*
 1059 *Entomol* 60:77-102
 1060
 1061 Scheffrahn RH, Scherer CW (2013) Efficacy of Altriset® on the tropical arboreal
 1062 termite, *Nasutitermes corniger* (Isoptera: Termitidae: Nasutitermitinae). *Fla Entomol*
 1063 96:249-251
 1064
 1065 Scheffrahn RH, Cabrera BJ, Kern Jr WH, Su N-Y (2002) *Nasutitermes costalis*
 1066 (Isoptera: Termitidae) in Florida: first record of a non-endemic establishment by a
 1067 higher termite. *Fla Entomol* 85:273-275
 1068
 1069 Scheffrahn RH, Hochmair HH, Kern Jr WH, Warner J, Krecek J, Maharajh B, Cabrera BJ,
 1070 Hickman R (2014) Targeted elimination of the exotic termite, *Nasutitermes corniger*
 1071 (Isoptera: Termitidae: Nasutitermitinae), from infested tracts in southeastern Florida.
 1072 *Int J Pest Manage* 60:9-21
 1073
 1074 Scheffrahn RH, Krecek J, Szalanski AL, Austin JW (2005a) Synonymy of neotropical
 1075 arboreal termites *Nasutitermes corniger* and *N. costalis* (Isoptera: Termitidae:
 1076 Nasutitermitinae), with evidence from morphology, genetics, and biogeography. *Ann*
 1077 *Entomol Soc Am* 98:273-281
 1078
 1079 Scheffrahn RH, Krecek J, Szalanski AL, Austin JW, Roisin Y (2005b) Synonymy of two
 1080 arboreal termites (Isoptera: Termitidae: Nasutitermitinae): *Nasutitermes corniger* from
 1081 the Neotropics and *N. polygynus* from New Guinea. *Fla Entomol* 88:28-33
 1082

1083 Scheffrahn RH, Su N-Y, Diehl B (1990) Native, introduced, and structure-infesting
 1084 termites of the Turks and Caicos Islands, BWI (Isoptera: Kalotermitidae,
 1085 Rhinotermitidae, Termitidae). Fla Entomol 73:622-627
 1086

1087 Schultz TP, Nicholas DD, McIntyre CR (2008) Recent patents and developments in
 1088 biocidal wood protection systems for exterior applications. Recent Patents on
 1089 Materials Science 1:128-134
 1090

1091 Schultz TP, Nicholas DD, Preston AF (2007) A brief review of the past, present and
 1092 future of wood preservation. Pest Manage Sci 63:784-788
 1093

1094 Serit M, Ishida M, Hagiwara N, Kim M, Yamamoto T and Takahashi S (1992) Meliaceae
 1095 and rutaceae limonoids as termite antifeedants evaluated using *Reticulitermes speratus*
 1096 kolbe (Isoptera: Rhinotermitidae). J Chem Ecol 18:593-603
 1097

1098 Sharma S, Vasudevan P, Madan M (1991) Insecticidal value of castor (*Ricinus cummunis*)
 1099 against termites. Int Biodeter Biodegr 27:249-254
 1100

1101 Sharma S, Verma M, Prasad R, Yadav D (2011) Efficacy of non-edible oil seedcakes
 1102 against termite (*Odontotermes obesus*). J Sci Ind Res 70:1037-1041
 1103

1104 Shelton TG, Grace K (2003) Termite Physiology in relation to wood degradation and
 1105 termite control. In: Goodell B, Nicholas DD, Schultz TP, Wood deterioration and
 1106 preservation: advances in our changing world, San Diego, California, pp. 242-250
 1107

1108 Siderhurst MS, James DM, Blunt TD, Bjostad LB (2005) Antimicrobial activity of the
 1109 termite (Isoptera) alkaloid norharmane against the entomopathogenic fungus
 1110 *Metarhizium anisopliae*. Sociobiology 46:563-577
 1111

1112 Sillam-Dussès D, Sémon E, Robert A, Cancellato E, Lenz M, Valterova I, Bordereau C
 1113 (2010) Identification of multi component trail pheromones in the most evolutionarily
 1114 derived termites, the Nasutitermitinae (Termitidae). Biol J Linn Soc 99:20-27
 1115

1116 Silva MD, Sá RA, Napoleão TH, Gomes FS, Santos ND, Albuquerque AC, Xavier HS,
 1117 Paiva PM, Correia MT, Coelho LC (2009) Purified *Cladonia verticillaris* lichen lectin:
 1118 Insecticidal activity on *Nasutitermes corniger* (Isoptera: Termitidae). Int Biodeter
 1119 Biodegr 63:334-340
 1120

1121 Silvestri, F. 1903. Contribuzione alla conoscenza dei Termitidi e Termitofili dell’America
 1122 Meridionale. Redia 1: 1–234
 1123

1124 Smeathman, H. 1781. Some account of the termites, which are found in Africa and other
 1125 hot climates. Philos Trans R Soc Lond 71:139–192
 1126

1127 Snyder, T.E. 1949. Catalog of the termites (Isoptera) of the World. Smithsonian
 1128 Miscellaneous Collections 112:1–490
 1129

1130 Souza JD, Silva MB, Argolo AC, Napoleão TH, Sá RA, Correia MT, Paiva PM, Silva
 1131 MD, Coelho LC (2011) A new *Bauhinia monandra* galactose-specific lectin purified

1132 in milligram quantities from secondary roots with antifungal and termiticidal
 1133 activities. Int Biodeter Biodegr 65:696-702
 1134

1135 Stuart AM (1963) Origin of the trail in the termites *Nasutitermes corniger* (Motschulsky)
 1136 and *Zootermopsis nevadensis* (Hagen), Isoptera. Physiol Zool 36:69-84
 1137

1138 Su N-Y and Scheffrahn RH (1998) A review of subterranean termite control practices and
 1139 prospects for integrated pest management programs. Integrated Pest Management
 1140 Review 3:1-13
 1141

1142 Su N-Y and Scheffrahn RH (1993) Laboratory evaluation of two chitin synthesis
 1143 inhibitors, hexaflumuron and diflubenzuron, as bait toxicants against the Formosan
 1144 subterranean termite (Isoptera: Rhinotermitidae). J Econ Entomol 86:1453-1457
 1145

1146 Tahiri A, Amissa adima A, Adje F, Amusant N (2011) Effet pesticide et screening des
 1147 extraits de *Azadirachta indica* (A.) Juss. sur le termite *Macrotermes bellicosus*
 1148 Rambur. Bois For Trop 310:79-88
 1149

1150 Tāyasu I, Sugimoto A, Wada E, Abe T (1994) Xylophagous termites depending on
 1151 atmospheric nitrogen. Naturwissenschaften 81:229-231
 1152

1153 Taylor AM, Gartner BL, Morrell JJ (2002) Heartwood formation and natural durability –
 1154 a review. Wood Fiber Sci 34:587-611
 1155

1156 Thorne BL (1983) Alate production and sex ratio in colonies of the Neotropical termite
1157 *Nasutitermes corniger* (Isoptera; Termitidae). *Oecologia* 58:103-109
1158
1159 Thorne BL (1980) Differences in nest architecture between the neotropical arboreal
1160 termites, *Nasutitermes corniger* and *Nasutitermes ephratae* (Isoptera: Termitidae).
1161 *Psyche* 87:235-244
1162
1163 Thorne BL (1984) Polygyny in the Neotropical termite *Nasutitermes corniger*: life history
1164 consequences of queen mutualism. *Behav Ecol Sociobiol* 14:117-136
1165
1166 Thorne BL, Collins MS, Bjorndal KA (1996) Architecture and nutrient analysis of
1167 arboreal carton nests of two neotropical *Nasutitermes* species (Isoptera: Termitidae),
1168 with notes on embedded nodules. *Fla Entomol* 79:27
1169
1170 Traniello JF, Busher C (1985) Chemical regulation of polyethism during foraging in
1171 the neotropical termite *Nasutitermes costalis*. *J Chem Ecol* 11:319-332
1172
1173 Traniello JF (1981) Enemy deterrence in the recruitment strategy of a termite: soldier-
1174 organized foraging in *Nasutitermes costalis*. *Proc Nat Acad Sci* 78:1976-1979
1175
1176 UNEP/FAO (2000) Global IPM Facility Expert Group on Termite Biology and
1177 Management. Finding alternatives to persistent organic pollutants (POPs) for termite
1178 management.<http://www.unep.org/chemicalsandwaste/Portals/9/Pesticides/Alternative>
1179 [s-termite-fulldocument.pdf](http://www.unep.org/chemicalsandwaste/Portals/9/Pesticides/Alternative)
1180

- Vasconcellos A and Moura FMDS (2010) Wood litter consumption by three species of *Nasutitermes* termites in an area of the Atlantic Coastal Forest in Northeastern Brazil. *J Insect Sci* 10:72
- Verma M, Sharma S, Prasad R (2009) Biological alternatives for termite control: a review. *Int Biodeter Biodegr* 63:959-972
- Warnecke F, Luginbühl P, Ivanova N, Ghassemian M, Richardson TH, Stege JT, Cayouette M, McHardy AC, Djordjevic G, Aboushadi N, Sorek R, Tringe SG, Podar M, Martin HG, Kunin V, Dalevi D, Madejska J, Kirton E, Platt D, Szeto E, Salamov A, Barry K, Mikhailova N, Kyrpides NC, Matson EG, Ottesen EA, Zhang X, Hernandez M, Murillo C, Acosta LG, Rigoutsos I, Tamayo G, Green BD, Chang B, Rubin EM, Mathur EJ, Robertson DE, Hugenholtz P, Leadbetter JR (2007) Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. *Nature* 450:560-565
- YunLing K, GuanJun C, ShuAng Z, TianYong Z, ChunXiao W, WeiJin T (2009) Bioactivities of azadirachtin against *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Sociobiology* 53:677-685
- Zhao C, Rickards RW, Trowell SC (2004) Antibiotics from Australian terrestrial invertebrates. Part 1: Antibacterial trinervitadienes from the termite *Nasutitermes triodiae*. *Tetrahedron* 60:10753-10759

Figure 1: Cumulative number of valid species of *Nasutitermes* since 1758.




Figure 2: Geographical distribution of *Nasutitermes corniger*. This termite is native of Central and South America and the Caribbean islands, and is invasive in the Bahamas, Florida and New Guinea (according to Scheffrahn et al. and Evans et al.).^{7,9}






Figure 3: *Nasutitermes corniger*'s colony with the king (A), queen (B), eggs (C), alate (D), adultoid (E), ergatoid (F), large (G1) and small (G2) workers, soldier (H), and gut's microbiome (photo according to Mikaelyan et al. 2014) (I).

Dotted arrows indicate symbiotic relationships. Gray circles indicate targets of all pest management strategies found in the literature.

Figure 4: Soldier (left) and major worker (right) of *Nasutitermes corniger*, same magnification. © Copyright 2016 R. Constantino.




Figure 5: Arboreal nest of *Nasutitermes corniger* on a cacao tree. © Copyright 2016 R. Constantino.



Table 1: Examples of the synthetic chemical products used for termite management.

Classes	Active compounds	References
Arsenic derivatives	Chromated copper arsenate (CCA)	Schultz et al. 2007
	Arsenic trioxide	
	Calcium arsenate	Verma et al. 2009; UNEP 2000; Santana et al. 2010
Copper derivatives	Alkaline copper quat (ACQ)	UNEP 2000; Schultz et al. 2007
	Copper azole	
	Copper naphthenate	
Zinc derivatives	Zinc naphthenate	UNEP 2000; Schultz et al. 2007
Borate formulations	Disodium octaborate tetrahydrate (DOT)	UNEP 2000; Schultz et al. 2007; Clausen and Yang 2007
Pyrethroids	Permethrin	UNEP 2000; Schultz et al. 2007
Neonicotinoids	Imidacloprid	Schultz et al. 2007;
Phenylpyrazoles	Thiamethoxan	Scheffrahn et al. 2014
	Fipronil,	Verma et al. 2009; UNEP 2000; Scheffrahn et al. 2014, Santana et al. 2010
Pyrethroid ethers	Chlorfenapyr	Verma et al. 2009; UNEP 2000; Santana et al. 2010
	Silafluofen	Verma et al. 2009; UNEP 2000; Santana et al. 2010
Organophosphate compounds	Chlorpyrifos	Verma et al. 2009; UNEP 2000; Santana et al. 2010
	Fenitrothion	
	Phenthoate	
	Triazophos	
Organochlorate compounds	Chlorothalonil	Verma et al. 2009; UNEP 2000; Santana et al. 2010
Benzoylureas	Chlorfluazuron	Verma et al. 2009; UNEP 2000; Santana et al. 2010
	Diffubenzuron	
	Hexaflumuron	
	Triflumuron	
Anthralinic diamides	Chlorantraniliprole	Verma et al. 2009; UNEP 2000; Santana et al. 2010; Scheffrahn and Scherer 2013
Aminohydrazones	Hydramethylnon	Verma et al. 2009; UNEP 2000; Santana et al. 2010
Spinosoids	Spinosad	Verma et al. 2009; UNEP 2000; Santana et al. 2010
Sulfonamides	Sulfluramid	Verma et al. 2009; UNEP 2000; Santana et al. 2010
Organic and inorganic gas	Carbon dioxide	Verma et al. 2009; UNEP 2000; Santana et al. 2010; Scheffrahn et al. 2014
	Methyl bromide	
	Phosphine	
	Sulfuryl fluoride	

Table 2: Sustainable management with antitermitic and antimicrobial botanicals and with entomopathogens.

Termite	Plant (Family)/part of plant or Entomopathogens (Family)	Activity	Active extracts / compounds	References
Antitermitic botanicals				
<i>N. corniger</i>	<i>Diospyros dichroa</i> Sandw. (Ebenaceae)	Repellent activity (> 90%) at 3.2 µg/mm ² (40 mg/ml)	Wood ethyl acetate extract	Pers. results based on Ganapaty et al 2004, Rodrigues 2010, and Boulogne et al 2011 previous works
<i>N. corniger</i>	<i>Lonchocarpus chrysophyllus</i> Kleinhoonte (Fabaceae)	Repellent activity (> 90%) at 3.2 µg/mm ² (40 mg/ml)	Bark ethyl acetate and petroleum ether extracts	Pers. results (Boulogne et al. <i>in prep</i>)
<i>N. corniger</i>	<i>Myracrodruon urundeuva</i> Allemão (Anacardiaceae)/ heartwood	No repellent activity Contact toxicity (LC ₅₀ =0.248 mg/mL for workers and 0.199 mg/mL for soldiers)	Lectins	Sá et al. 2008
<i>N. corniger</i>	<i>Myracrodruon urundeuva</i> Allemão (Anacardiaceae)/ heartwood and bark	No contact toxicity Repellent activity	Crude methanolic extract	Sá et al. 2009
<i>N. corniger</i>	<i>Anadenanthera colubrina</i> (Vell.) Brenan (Fabaceae)/ heartwood	Contact toxicity (LC ₅₀ = 7.3 mg/mL for ethyl acetate extract, LC ₅₀ = 67.4 mg/mL for cyclohexane extract after 4 days)	Cyclohexane and ethyl acetate extracts	Santana et al. 2010
<i>N. corniger</i>	<i>Hymenaea stigonocarpa</i> Hayne (Fabaceae)/ heartwood	Contact toxicity (LC ₅₀ =11.9 mg/mL after 4 days)	Cyclohexane extract	Santana et al. 2010
<i>N. corniger</i>	<i>Bowdichia virgilioides</i> Kunth, (Fabaceae)/ heartwood	Contact toxicity (LC ₅₀ =7.2 mg/mL after 4 days)	Alkaloidal extract	Santana et al. 2010
<i>N. corniger</i>	<i>Bowdichia virgilioides</i> Kunth, (Fabaceae)/ heartwood	Toxic (LC ₅₀ =12.2 mg/mL after 4 days) and repellent activities	Cyclohexane extract	Santana et al. 2010
<i>N. macrocephalus</i>	<i>Sextonia rubra</i> (Mez) Van der Werff (Lauraceae)/ heartwood	Contact toxicity (LC ₁₀₀ <1.9 µg/mm ² after 72h)	Ethyl-acetate extract/ rubrynolide	Rodrigues et al. 2011
<i>Nasutitermes</i> sp	<i>Bagassa guianensis</i> Aubl. (Moraceae), <i>Erisma uncinatum</i> Warm. (Vochysiaceae), <i>Buchenavia parvifolia</i> Ducke (Combretaceae), <i>Dinizia excelsa</i> Ducke (Fabaceae) <i>Pouteria guianensis</i> Aubl. (Sapotaceae)/ wood	Wood impregnation conferred durability to <i>Cecropia</i> sp. (Urticaceae) and <i>Simarouba amara</i> Aubl. (Simaroubaceae) woods		Barbosa et al. 2007; Peres Filho et al. 2006
<i>N. macrocephalus</i>	<i>Diospyros dichroa</i> Sandw. (Ebenaceae)/ wood	Contact toxicity (LC ₁₀₀ <1.9 µg/mm ² and 5.6 µg/mm ² after 72h)	Methanolic extract and Ethyl acetate extract.	Harun and Labosky 1985; Rodrigues 2010
<i>N. corniger</i>	<i>Bauhinia monandra</i> Kurz (Fabaceae)/ roots	Contact toxicity (LC ₅₀ = 0.09 mg/mL for workers and 0.395 for soldiers after 12 days)	Lectins	Souza et al. 2011
<i>N. corniger</i>	<i>Opuntia ficus-indica</i> (L.) Mill. (Cactaceae)/ cladodes	Contact toxicity (LC ₅₀ =0.116 mg/mL for workers and 1.5 for soldiers)	Lectins	Paiva et al. 2011
<i>N. corniger</i>	<i>Moringa oleifera</i> Lam. (Moringaceae)/ seeds	No repellent activity Contact toxicity at 1.0 and 1.5 mg/mL	Lectins	Paiva et al. 2011
<i>N. corniger</i>	<i>Myracrodruon urundeuva</i> Allemão (Anacardiaceae)/ leaves	Contact toxicity (LC ₅₀ =0.374 to 0.974 mg/ml on workers and 0.432 to 0.787 mg/ml on soldiers)	Lectins	Napoleão et al. 2011
<i>N. corniger</i>	<i>Cladonia verticillaris</i>	Contact toxicity	Lectins	Silva et al. 2009

	(Cladoniaceae) / lichen	(LC ₅₀ = 0.196 mg/mL on workers and 0.5 mg/mL on soldiers after 10 days)		
<i>N. corniger</i>	<i>Microgramma vaccinifolia</i> (Langsd. & Fisch.) Copel. (Polypodiaceae)/ rhizome	Contact toxicity Ability to disrupt trypsin-like protease, acid phosphatase, and cellulase activities in termite's gut	Lectins	Lima et al. 2014
<i>Coptotermes formosanus</i> , <i>Coptotermes gestroi</i> , <i>Reticulitermes speratus</i> , <i>Macrotermes bellicosus</i> , <i>Macrotermes nigeriensis</i> , <i>Odontotermes obes</i>	<i>Azadirachta indica</i> A. Juss. (Meliaceae) <i>Ricinus communis</i> L. (Euphorbiaceae)	Contact toxicity , metabolic poison, repellent or antifeedant activities		Sharma et al. 1991; Grace and Yates 1992; Serit et al. 1992; Ivbijaro et al. 1993; Delate and Grace 1995; Chuanchuan et al. 1998; UNEP 2000; Yunling et al. 2009; Sharma et al. 2011; Tahiri et al. 2011; Himmi et al. 2013
Antimicrobial botanicals				
<i>N. corniger</i>	<i>Myracrodruon urundeuva</i> Allemão (Anacardiaceae)/ bark, heartwood and leaf	Antibacterial activity on gut's symbiotic bacteria (MIC from 62.5 to 250 mg/mL)	Lectins	Napoleão et al. 2011
<i>Microtermes obesi</i>	<i>Withania somnifera</i> (L.) Dunal (Solanaceae), <i>Croton tiglium</i> L. (Euphorbiaceae) and <i>Hygrophila auriculata</i> Schumach.) Heine (Acanthaceae) / seeds and leaves	Reduction of colonies number of bacterial symbionts		Ahmed et al. 2006
<i>Coptotermes formosanus</i>	<i>Chrysopogon zizanioides</i> (L.) Roberty (syn. <i>Vetiveria zizanioides</i> (L.) Nash) (Poaceae)	Reduction number of hindgut flagellate species (more specially <i>Pseudotrichonympha grassii</i> , an important cellulolytic flagellate protist)	Oil/ Nootkatone	Maistrello et al. 2003; 2011
<i>Coptotermes formosanus</i>	<i>Azadirachta indica</i> A. Juss. (Meliaceae)	Reduction of population of <i>P. grassii</i> and <i>Spirochaetes</i>	Azadirachtin	Doolittle et al. 2007
Entomopatogens				
<i>Nasutitermes</i> sp.	<i>Termitaria coronata</i> (Deuteromycetes, Termitariales)			Ensaf et al. 2006
<i>N. corniger</i> and <i>Nasutitermes</i> sp.	<i>Metarhizium</i> sp			Chouvenec et al. 2011; Salas-Acuña 2012