

**A diet containing soybean oil heated for three hours
increases adipose tissue weight but decreases body
weight in C57BL/6 J mice**

Meera Penumetcha, Mary K Schneider, Holly A Cheek, Sonia Karabina

► **To cite this version:**

Meera Penumetcha, Mary K Schneider, Holly A Cheek, Sonia Karabina. A diet containing soybean oil heated for three hours increases adipose tissue weight but decreases body weight in C57BL/6 J mice. *Lipids in Health and Disease*, BioMed Central, 2013, 12, pp.26. <10.1186/1476-511X-12-26>. <hal-01583973>

HAL Id: hal-01583973

<http://hal.upmc.fr/hal-01583973>

Submitted on 8 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



SHORT REPORT

Open Access

A diet containing soybean oil heated for three hours increases adipose tissue weight but decreases body weight in C57BL/6 J mice

Meera Penumetcha^{1*}, Mary K Schneider¹, Holly A Cheek¹ and Sonia Karabina²

Abstract

Background: Our previous work showed that dietary oxidized linoleic acid given, as a single fatty acid, to LDL receptor knockout mice decreased weight gain as compared to control mice. Other studies have also reported that animals fed oils heated for 24 h or greater showed reduced weight gain. These observations, while important, have limited significance since fried foods in the typical human diet do not contain the extreme levels of oxidized lipids used in these studies. The main goal of this study was to investigate the effects of a diet containing soybean oil heated for 3 h on weight gain and fat pad mass in mice. Additionally, because PPAR γ and UCP-1 mediate adipocyte differentiation and thermogenesis, respectively, the effect of this diet on these proteins was also examined.

Findings: Four to six week old male C57BL/6 J mice were randomly divided into three groups and given either a low fat diet with heated soybean oil (HSO) or unheated soybean oil (USO) or pair fed for 16 weeks. Weight and food intake were monitored and fat pads were harvested upon the study's termination. Mice consuming the HSO diet had significantly increased fat pad mass but gained less weight as compared to mice in the USO group despite a similar caloric intake and similar levels of PPAR γ and UCP1.

Conclusion: This is the first study to show that a diet containing soybean oil heated for a short time increases fat mass despite a decreased weight gain in C57BL/6 J mice. The subsequent metabolic consequences of this increased fat mass merits further investigation.

Keywords: Heated oil, Oxidized oil, Adipose mass, Diet

Findings

Background

Foods rich in polyunsaturated fatty acids (PUFA) are susceptible to oxidation. Food processing, storage and cooking at high temperatures hasten this process and result in the formation of both primary (peroxide value, PV; conjugated dienes, CD) and secondary products of lipid oxidation. Although studies that investigate the biological effects of these dietary oxidized lipids (DOL) in humans are limited, rodents that are fed diets containing large proportions (50-100% of total fat) of oils heated for prolonged periods (10 h to 38 d) gain less weight and show a reduction in fat mass as compared to controls

[1-3]. A literature review of commonly consumed foods, however, demonstrates that French fries, fried fish, pizza and nuts have PVs of 1.98, 2.2, 2.45 and 1.20 meq/Kg, respectively [4], which are significantly lower than those used in previous animal studies. We thus believe that these studies do not accurately represent the effects of typical DOL consumption in the human population. From the above mentioned studies it appears that the weight reducing effects of DOL are probably due to the high amounts of oxidized lipids used. In order to address this issue, we used a diet containing oil heated for only 3 h, comprising about 1/3 of total dietary fat (wt/wt), and tested its effects on total body weight and fat pad mass. As consumption of fried and processed foods is increasing globally it is important to know how DOL influence adipose tissue mass and subsequent metabolic risk.

* Correspondence: mpenumetcha@gsu.edu

¹Division of Nutrition, Byrdine F. Lewis School of Nursing and Health Professions, Georgia State University, Paris Cedex 12, France
Full list of author information is available at the end of the article

Methods

Animals

Twenty-four, 6–8 week old, male C57BL/6 J mice were purchased from the Jackson Laboratory (Bar Harbor, ME). The Georgia State University (GSU) Animal Care Committee approved all protocols (protocol #A060654) based on the Helsinki Declaration and mice were treated in compliance with the GSU Animal Care Committee regulations. Mice were randomly divided into three groups. One group was fed, *ad libitum*, a low fat diet with one-third of the total fat provided by unheated soybean oil (USO) while a second group was fed *ad libitum* a low fat diet with one-third of the total fat provided by heated soybean oil (HSO). A third group of mice was pair-fed (PSO) to the HSO group. In other words, though the PSO mice were fed the diet containing unheated oil, they were only allowed to consume the same amount of food (in grams) as the HSO group. A known amount of fresh food was placed in the cage every two days and any remaining or spilled food was weighed and discarded. Mice were weighed once weekly on a digital scale. After 16 weeks, mice were euthanized following an overnight fast. An incision was made in the abdomen and epididymal white adipose tissue (EWAT) was removed. The mouse was then placed on its ventral side and the interscapular brown adipose tissue (IBAT) was carefully excised. Fat pads were quickly rinsed in ice cold PBS, weighed on a high precision balance, flash frozen in liquid nitrogen and stored at -80°C . Methods for Western blot and RT-QPCR are reported in Additional file 1.

Mouse diet

HSO diet was produced by heating soybean oil on a hot plate at 190°C for 3 hours with an air compressor providing a continuous flow of oxygen throughout the oil. Both heated and unheated soybean oils were sent to Research Diets Inc. (New Brunswick, NJ) on dry ice to be incorporated into pelleted mouse chow. Each diet was distinctively colored to facilitate observation of fecal samples. Mouse diet was stored at 4°C to prevent further oxidation.

PV of the mouse chow was determined by CII Laboratory Services (Kansas City, MO). Lipids were extracted

from all diets using a solvent system of hexane and isopropanol (HIP) (3:2) as described by Radin [5]. Conjugated dienes and TBARS (secondary product of oxidation) of the extracted lipids were determined as described previously [6]. Compositions of the diets along with the values for products of oxidation are listed in Additional file 2: Table S2.

Statistical analysis

Statistics were generated using SPSS version 16.0 (SPSS Inc., Chicago, IL) using either one way-ANOVA followed by Bonferroni Post hoc tests or Kruskal-Wallis test followed by Mann Whitney test depending on the distribution. Statistical significance was set at $P \leq 0.05$.

Results

Food intake among the USO, HSO and PSO groups was not different (Table 1). However, mice fed the USO diet gained the most weight in comparison to mice that were pair fed or consumed the HSO diet (Table 1) Mean body weights over time are shown in Additional file 3: Figure S2. Food efficiency was similar in the USO and HSO groups but was significantly lower in the PSO group (Table 1). Despite the lesser weight gain in the HSO group as compared to the USO group, EWAT and IBAT weights were significantly higher in the HSO group compared to the USO and PSO groups (Figure 1a). EWAT PPAR γ mRNA was not different between the groups (Figure 1b). PPAR γ and UCP1 mRNA levels in IBAT were not different between the three groups (Figure 1b). PPAR γ and UCP-1 protein expression levels were not different between the groups in both EWAT and IBAT (Table 1).

Discussion

We successfully created a mouse diet that contained mostly primary products of oxidation (PV and CD) and negligible amounts of secondary products of oxidation (TBARS). In this study, mice that consumed a diet containing 1/3 of the total fat from heated oil gained less weight as compared to mice that consumed the unheated oil diet. This decrease in weight gain was 6% in our study as compared to 15-55% in other studies [1-3,7]. In those studies, oils were highly oxidized and contained 150 times

Table 1 Body weight gain, food intake and other parameters (mean \pm SD) in mice fed a USO, HSO or PSO diet

Parameter	USO	HSO	PSO
Food intake (g/wk)	20.65 \pm 0.09 ^a	18.4 \pm 0.05 ^a	18.38 \pm 0.19 ^a
Weight gain (g)	8.86 \pm 1.37 ^a	7.10 \pm 1.47 ^b	5.71 \pm 1.13 ^c
Food efficiency (weight gain, g/food intake, g)	0.031 \pm 0.005 ^a	0.028 \pm 0.006 ^{ab}	0.022 \pm 0.004 ^b
EWAT PPAR γ protein (PPAR γ /GAPDH)	0.226 \pm 0.08 ^a	0.264 \pm 0.12 ^a	0.456 \pm 0.6 ^a
IBAT UCP-1 protein (UCP-1/Sypro protein stain)	0.139 \pm 0.16 ^a	0.188 \pm 0.16 ^a	0.127 \pm 0.20 ^a

USO=Unheated soybean oil diet; HSO=Heated soybean oil diet; PSO=Pair-fed to HSO with USO. Differences between the three groups were compared with one-way ANOVA followed by Bonferroni pair wise comparisons. Mean values within a row, with different superscripts of the alphabet, are significantly different from one another. N=8 per group for all the parameters except for food intake. Mice were housed at 2-3/cage and so n=3 for food intake ($p \leq 0.05$).




Figure 1 Mean fat pad weights (1a) and mRNA expression (1b) of mice fed a USO (white bar; n=8), HSO (black bar; n=8) or PSO (grey bar; n=8) diet. Bars represent mean \pm SD. Group means were compared by the Kruskal Wallis test and pair wise comparisons were made by Mann Whitney U test. Statistical significance was set at $p \leq 0.05$. Means with “†” are significantly different as compared to USO and PSO groups.

the PV and 1500 times the TBARS as compared to the diet in this study. Oxidized oils have been shown to decrease fat absorption [8] and increase fecal fat excretion. Feces from mice across all groups looked similar and were *not* oily. Thus, the lower weight gain in the HSO group could be due to factors other than decreased fat absorption. Surprisingly, pair-fed mice (PSO) gained less weight than the HSO group, despite a similar caloric intake and physical activity (data not shown). Pair feeding, however, required that only a small amount of food (3–4 pellets) be placed in the feeding hopper and this might have forced the mice to exert more energy to access the food, resulting in increased energy expenditure.

A novel finding in our study is the increase in fat pad mass seen in mice consuming mildly heated soybean oil as compared to mice in the USO and PSO groups. Studies by Chao et al., in mice [1] and by others in rats [9] have shown that diets containing heated oils decrease fat pad mass and body weight, respectively. As mentioned before, this could be reflective of the deficit in calories due to decreased fat absorption. The gain in fat mass in this study cannot be explained by increased food intake because there were no differences in food intake between the USO and HSO groups. One explanation for the increased fat pad weight among HSO as compared to USO mice may be the repartitioning of fat and lean tissue. For example, conjugated linoleic acid, which is very similar in structure to oxidized PUFA (hydroxy linoleic acid), has been shown to change the proportions of lean and fat mass in the absence of weight change [10]. In addition, primary products of lipid oxidation, but not secondary products, have been shown to act as ligands of PPAR γ [11]. Activation of PPAR γ promotes modest increases in fat mass as seen in mice [12] and humans [13] treated with rosiglitazone, a known PPAR γ agonist. We hypothesized that the increase in fat pad mass among mice consuming HSO could be due to an upregulation of PPAR γ . Both mRNA and protein levels of PPAR γ , however, were not different in the

EWAT and IBAT between the groups. This suggests that mildly heated soybean oil might not change PPAR γ mRNA or protein levels, but it cannot be ruled out that it could activate PPAR γ . Furthermore the large variance in measuring PPAR γ along with the small sample size might have made it difficult to find a statistically significant difference.

A limitation of this study is that energy expenditure was not measured in these mice and would have answered the question of whether oxidized lipids could alter basal metabolic rate thereby explaining the differences in weight gain. Thermogenesis is a component of energy balance. We wanted to see if the reduced weight gain in the HSO group could be attributed to an energy deficit and measured IBAT UCP-1, a recognized cellular marker of thermogenesis. As we report here, there were no differences in the UCP-1 mRNA or protein levels between any of the groups. The lack of difference in UCP-1 levels in the present study, however, does not negate a role for thermogenesis. Evidence from studies in mice show that changes in thermogenesis can occur independent of UCP-1 expression [14]. In addition, measuring thermogenesis by implanting a subcutaneous sensor would yield a direct measurement.

In conclusion, this is the first study to show that dietary oxidized lipids, in amounts found in commonly consumed foods, can increase fat pad mass. The metabolic consequences of this increased fat mass with regard to glucose homeostasis merits further investigation.

Additional files

Additional file 1: Methods. Western blots and RT-QPCR.

Additional file 2: Table S2. Composition of the HSO^a and USO^b diets.

Additional file 3: Figure S2. Mean body weights of mice over the 16 week study period.

Abbreviations

HSO: Heated soybean oil; PSO: Mice pair-fed with unheated soybean oil; EWAT: Epididymal white adipose tissue; IBAT: Interscapular brown adipose tissue; PPAR γ : Peroxisome proliferator activated receptor gamma; UCP1: Uncoupling protein 1; DOL: Dietary oxidized lipids; CD: Conjugated diene; PV: Peroxide value.

Competing interests

The authors declare that they have no competing interests to report.

Authors' contributions

MP designed the study, participated in animal work, analyzed and interpreted the data and drafted the manuscript. MKS participated in the weighing, feeding and surgical procedures of mice, Western blot analysis of PPAR γ and reviewed the manuscript. HAC carried out the Western blot analysis of UCP-1 and reviewed the manuscript. SK was responsible for carrying out the RT-QPCR, data analysis and reviewed the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank Dr. Timothy Bartness and his laboratory for their technical and scientific assistance in conducting this study. We would like to thank Welch, Holme & Clark Co (NJ, USA) for providing soybean oil free of charge.

Author details

¹Division of Nutrition, Byrdine F. Lewis School of Nursing and Health Professions, Georgia State University, Paris Cedex 12, France. ²INSERM UMRS 933, UPMC-Paris 6, 26, avenue Arnold Netter, 75571, Paris Cedex 12, France.

Received: 26 September 2012 Accepted: 28 February 2013

Published: 6 March 2013

References

1. Chao PM, Huang HL, Liao CH, Huang ST, Huang CJ: **A high oxidised frying oil content diet is less adipogenic, but induces glucose intolerance in rodents.** *Br J Nutr* 2007, **98**(1):63–71.
2. Liao CH, Shaw HM, Chao PM: **Impairment of glucose metabolism in mice induced by dietary oxidized frying oil is different from that induced by conjugated linoleic acid.** *Nutrition* 2008, **24**(7–8):744–752.
3. Sulzle A, Hirche F, Eder K: **Thermally oxidized dietary fat upregulates the expression of target genes of PPAR alpha in rat liver.** *J Nutr* 2004, **134**(6):1375–1383.
4. Arora P: *Consumption of oxidized lipid and its association with markers of adiposity: NHANES 2006–2006.* Atlanta: Georgia State University; 2011.
5. Hara A, Radin NS: **Lipid extraction of tissues with a low-toxicity solvent.** *Anal Biochem* 1978, **90**(1):420–426.
6. Khan-Merchant N, Penumetcha M, Meilhac O, Parthasarathy S: **Oxidized fatty acids promote atherosclerosis only in the presence of dietary cholesterol in low-density lipoprotein receptor knockout mice.** *J Nutr* 2002, **132**(11):3256–3262.
7. Chao PM, Chao CY, Lin FJ, Huang C: **Oxidized frying oil up-regulates hepatic acyl-CoA oxidase and cytochrome P450 4 A1 genes in rats and activates PPARalpha.** *J Nutr* 2001, **131**(12):3166–3174.
8. Burenjargal M, Totani N: **Effects of thermally processed oil on weight loss in rats.** *J Oleo Sci* 2008, **57**(9):463–470.
9. Totani N, Burenjargal M, Yawata M: **Effects of oil heated with gluten on weight-loss dieting. I.** *J Oleo Sci* 2008, **57**(6):321–326.
10. Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW: **Effect of conjugated linoleic acid on body composition in mice.** *Lipids* 1997, **32**(8):853–858.
11. Ollero M, Junaidi O, Zaman MM, Tzamelis I, Ferrando AA, Andersson C, Blanco PG, Bialecki E, Freedman SD: **Decreased expression of peroxisome proliferator activated receptor gamma in cfr-/- mice.** *J Cell Physiol* 2004, **200**(2):235–244.
12. Yang KJ, Noh JR, Kim YH, Gang GT, Hwang JH, Yang SJ, Yeom YI, Lee CH: **Differential modulatory effects of rosiglitazone and pioglitazone on white adipose tissue in db/db mice.** *Life Sci* 2010, **87**(13–14):405–410.
13. Larsen TM, Toubro S, Astrup A: **PPARgamma agonists in the treatment of type II diabetes: is increased fatness commensurate with long-term efficacy?** *Int J Obes Relat Metab Disord* 2003, **27**(2):147–161.
14. Oudart H, Groscolas R, Calgari C, Nibelink M, Leray C, Le Maho Y, Malan A: **Brown fat thermogenesis in rats fed high-fat diets enriched with n-3 polyunsaturated fatty acids.** *Int J Obes Relat Metab Disord* 1997, **21**(11):955–962.

doi:10.1186/1476-511X-12-26

Cite this article as: Penumetcha *et al.*: A diet containing soybean oil heated for three hours increases adipose tissue weight but decreases body weight in C57BL/6 J mice. *Lipids in Health and Disease* 2013 **12**:26.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

