

Resistance of *M. leprae* to Quinolones: A Question of Relativity?

Nicolas Veziris, Aurélie Chauffour, Sylvie Escolano, Sarah Henquet, Masanori Matsuoka, Vincent Jarlier, Alexandra Aubry

► **To cite this version:**

Nicolas Veziris, Aurélie Chauffour, Sylvie Escolano, Sarah Henquet, Masanori Matsuoka, et al.. Resistance of *M. leprae* to Quinolones: A Question of Relativity?. PLoS Neglected Tropical Diseases, Public Library of Science, 2013, 7 (11), pp.e2559. <10.1371/journal.pntd.0002559>. <hal-01614678>

HAL Id: hal-01614678

<http://hal.upmc.fr/hal-01614678>

Submitted on 11 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Resistance of *M. leprae* to Quinolones: A Question of Relativity?

Nicolas Veziris^{1,2,3}, Aurélie Chauffour¹, Sylvie Escolano⁴, Sarah Henquet³, Masanori Matsuoka⁵, Vincent Jarlier^{1,2,3}, Alexandra Aubry^{1,2,3*}

1 Université Pierre et Marie Curie-Paris 6, Paris, France, **2** Centre National de Référence de la Résistance des Mycobactéries aux Antituberculeux, Paris, France, **3** Groupe Hospitalier Pitié-Salpêtrière, Assistance Publique Hôpitaux de Paris, Laboratoire de bactériologie-hygiène, Paris, France, **4** Inserm, Centre for Research in Epidemiology and Population Health (CESP), U1018, Biostatistics, F-94807, Villejuif, France and Univ Paris-Sud, UMRS 1018, F-94807, Villejuif, France, **5** Leprosy Research Center, National Institute of Infectious Diseases, Higashimurayama-shi, Tokyo, Japan

Abstract

Multidrug resistant leprosy, defined as resistance to rifampin, dapsone and fluoroquinolones (FQ), has been described in *Mycobacterium leprae*. However, the *in vivo* impact of fluoroquinolone resistance, mainly mediated by mutations in DNA gyrase (GyrA₂GyrB₂), has not been precisely assessed. Our objective was to measure the impact of a DNA gyrase mutation whose implication in fluoroquinolone resistance has been previously demonstrated through biochemical studies, on the *in vivo* activity of 3 fluoroquinolones: ofloxacin, moxifloxacin and garenoxacin.

Methodology/Principal Findings: We used the proportional bactericidal method. 210 four-week-old immunodeficient female nude mice (NMRI-Foxn1^{nu}/Foxn1^{nu}) were inoculated in the left hind footpad with 0.03 ml of bacterial suspension containing 5×10^3 , 5×10^2 , 5×10^1 , and 5×10^0 *M. leprae* AFB organisms of strain Hoshizuka-4 which is a multidrug resistant strain harboring a GyrA A91V substitution. An additional subgroup of 10 mice was inoculated with 5×10^{-1} bacilli in the untreated control group. The day after inoculation, subgroups of mice were treated with a single dose of ofloxacin, moxifloxacin, garenoxacin or clarithromycin at 150 mg/kg dosing. 12 months later mice were sacrificed and *M. leprae* bacilli were numbered in the footpad. The results from the untreated control group indicated that the infective inoculum contained 23% of viable *M. leprae*. The results from the moxifloxacin and garenoxacin groups indicated that a single dose of these drugs reduced the percentage of viable *M. leprae* by 90%, similarly to the reduction observed after a single dose of the positive control drug clarithromycin. Conversely, ofloxacin was less active than clarithromycin.

Conclusion/Significance: DNA gyrase mutation is not always synonymous of lack of *in vivo* fluoroquinolone activity in *M. leprae*. As for *M. tuberculosis*, *in vivo* studies allow to measure residual antibiotic activity in case of target mutations in *M. leprae*.

Citation: Veziris N, Chauffour A, Escolano S, Henquet S, Matsuoka M, et al. (2013) Resistance of *M. leprae* to Quinolones: A Question of Relativity? PLoS Negl Trop Dis 7(11): e2559. doi:10.1371/journal.pntd.0002559

Editor: Gerd Pluschke, Swiss Tropical and Public Health Institute, Switzerland

Received: June 11, 2013; **Accepted:** October 11, 2013; **Published:** November 14, 2013

Copyright: © 2013 Veziris et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Funding support for the study on mouse footpad susceptibility (testing animal facilities, animal keeper and technician) was provided by repeated annual grants from the Association Française Raoul Follereau. This experiment was supported by Association Française Raoul Follereau and Ordre de Malte-France (specific Grant MALTALÉP 2009). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: alexandra.aubry@upmc.fr

Introduction

Mycobacterium leprae is responsible for leprosy, that the World Health Assembly decided, in 1991, to “eliminate as a public health problem” by the year 2000. But, though a decreasing number of new cases registered each year (~219,000) during the recent years [1,2], it is generally admitted that the goal of leprosy elimination is far from being reached [3]. Future projections of the global leprosy burden indicates that 5 million new cases would arise between 2000 and 2020, and that in 2020 there would be still 1 million people with WHO grade 2 disability due to leprosy.

Reports from Asian countries with a high leprosy prevalence estimated rates of resistance at 15–20% to dapsone (DDS) and 3–8% to rifampin [4,5]. Some studies estimated around 50% of resistance to DDS in relapsing cases [6,7]. Although the exact magnitude of resistance to these drugs is difficult to assess,

resistance in *M. leprae* is a concern particularly in relapsing multi-bacillary leprosy patients, by strongly reducing the possibilities of an effective treatment [1,8–10].

Quinolones are good candidates for the development of more powerful treatments of leprosy, as demonstrated for moxifloxacin which is the only drug other than rifampin to be consistently bactericidal against *M. leprae* in clinical trials [11]. Fluoroquinolones play a crucial role in the treatment for drug-resistant leprosy and single-lesion new cases [11], but some quinolone-resistant *M. leprae* strains have been described [12,13]. The mode of action of quinolones against *M. leprae* has been clearly identified and mechanisms of resistance have been investigated [4,5,12–17]. Substitutions within a highly conserved region of GyrA and possibly GyrB, which are subunits of the tetrameric DNA gyrase (A₂B₂), the so-called quinolone resistance-determining region (QRDR), are associated with the development of ofloxacin

Author Summary

Although there is efficient multidrug therapy to cure leprosy, the transmission of *M. leprae* is still active, leading to 219,000 new cases in 2011. Drug resistant leprosy has been described and may prevent eradication of the disease, notably multidrug resistant defined as resistance to rifampin, dapsone and fluoroquinolones (FQ). Resistance to FQ is due to mutations in DNA gyrase. We used a mouse model to measure the impact of DNA gyrase mutations on *in vivo* FQ activity. All the FQ tested showed *in vivo* activity against the mutant tested (A91V mutant in subunit A of DNA gyrase). However, whereas ofloxacin was less active than the control treatment clarithromycin, it appeared that latter generation fluoroquinolones moxifloxacin and garenoxacin were as active as clarithromycin. Our results demonstrate that DNA gyrase mutation is not synonymous of total lack of *in vivo* FQ activity against *M. leprae*. Therefore, as for *M. tuberculosis*, *in vivo* studies are mandatory in order to measure the impact of DNA gyrase mutations on treatment efficacy against *M. leprae*.

resistance in *M. leprae*, as demonstrated before in *M. tuberculosis*. Due to the lack of *M. leprae* growth *in vitro*, the exact impact of DNA gyrase mutations on fluoroquinolone susceptibility remains largely unknown. We previously demonstrated, using an enzymatic assay, that GyrA substitutions do not have the same impact on all the fluoroquinolones [15]. As an example, garenoxacin had the same inhibitory activity against *M. leprae* DNA gyrase carrying mutation implicated in resistance to ofloxacin as against wild-type enzyme, underscoring the potential advantage of this compound in leprosy [15].

The aim of the present study was to evaluate the bactericidal activities of several quinolones (i.e. ofloxacin, moxifloxacin and garenoxacin) against a *M. leprae* strain carrying an A91V GyrA substitution known to be involved in quinolone resistance [18]. Relation between GyrA A91V mutation and ofloxacin resistance has been extensively proven in the literature both in patients experiencing relapse after ofloxacin treatment (notably for strain for the strain used in the present work) [12,18,19] and through DNA gyrase inhibition *in vitro* [15]. We demonstrated that despite GyrA A91V mutation, garenoxacin and moxifloxacin maintained some *in vivo* activity.

Materials and Methods

Ethics statement

The laboratory has been approved on April, 24th 2012 to carry out animal experiments. Nicolas Veziris who carried the animal experiments has the following license number: 75-1531. Aurélie Chauffour who performed the animal experiments has the following license number: B-75-1214. We followed the animal experiment guidelines of the Faculté de Médecine Pierre-et-Marie Curie. Animal experiments were performed in accordance with prevailing regulations regarding the care and use of laboratory animals by the European Commission. The experimental protocol was approved by the Departmental Direction of Veterinary services in Paris, France.

Materials

Hoshizuka-4 strain is a multidrug resistant strain with mutation in the three main genes involved in *M. leprae* resistance to antibiotics: *folP* gene (P55S), involved in dapsone resistance; *rpoB* gene (S456L), involved in rifampicin resistance; and *gyrA* gene

(A91V) involved in quinolones resistance (using numbering system of the *M. leprae* genome TN, GenBank nuNC002677). Briefly, Hoshizuka-4 strain was isolated from a patient who developed a lepromatous leprosy after repeated clinical relapses [18]. He received dapsone, streptomycin, rifampin, clofazimine, isoniazid, ofloxacin and prothionamide to treat subsequent relapses. These drugs were administered irregularly as monotherapy or in combinations, often at doses below recommended levels and standard multidrug therapy was never applied. The drug resistant profile of the isolated strain was confirmed by the mouse footpad method (in nude mice) for fluoroquinolones (two fluoroquinolones, ofloxacin and sparflaxacin, were tested at 2 concentrations, 0.0001% and 0.001%) [18]. The GyrA A91V substitution corresponds to amino acid 90 and 83 in *M. tuberculosis* and *E. coli* numbering system, respectively.

Four week-old Nude mice (NMRI-*Foxn1*^{nu}/*Foxn1*^{nu}) were purchased from JANVIER breeding center, Le Genest Saint-Isle, France.

Ofloxacin was purchased from Sanofi-Aventis, France; moxifloxacin from Bayer Santé, France; garenoxacin from EasyBuyer LTD, China, and clarithromycin from Abbot France, France.

Infection of mice with *M. leprae*

Animal experiments were performed in accordance with prevailing regulations regarding the care and use of laboratory animals by the European Commission. The experimental protocol was approved by the Departmental Direction of Veterinary services in Paris, France.

The 'proportional bactericidal' technique, described by Colston, allows measuring the bactericidal activity of a compound [20]. Mice are inoculated with serial 10-fold dilutions of the suspension of *M. leprae*. A group of mice were left untreated; the other mice are treated for a period of time that varies from a single dose to 60 days (usually 10 mice per dilution of inoculum for each treatment-group). After treatment, the mice are held for 12 months, to permit a single surviving organism to multiply to a readily detectable level (*M. leprae* divides every 14 days). Harvests of *M. leprae* are then performed from individual feet; the organisms are considered to have multiplied in those feet found to contain $\geq 10^5$ AFB. The proportion of viable *M. leprae* surviving treatment may then be calculated from the number of organisms that infects 50% of the mice. The proportion of viable *M. leprae* killed by the treatment is calculated by comparing the proportion of viable organisms in the treated mice to that in the control mice.

Two hundred and ten 4 week-old immunodeficient female Nude were divided among 5 groups, each containing four subgroups with 10 mice each. The mice of each subgroup were inoculated in the left hind footpad with 0.03 ml of bacterial suspension containing 5×10^3 , 5×10^2 , 5×10^1 , and 5×10^0 *M. leprae* organisms of strain Hoshizuka-4 [18]. The suspension needed to inoculate mice was prepared from one footpad harvested from a nude mouse (according to the Shepard and Mac Rae method [21]), that had been inoculated one year earlier in the lab with 6.10^4 AFB/footpad. Ten μ l of the suspension were taken to create slides and AFB/footpad were counted after Ziehl-Neelsen coloration and ten-fold dilutions were made if it was necessary. All further ten-fold dilutions were made in Hank's balanced salt. A fifth subgroup of the untreated control group was inoculated with 5×10^{-1} Acid Fast Bacilli (AFB) per footpad.

Treatment of mice

The day after inoculation, a table of randomization was created on website randomization.com in order to randomly allocate mice in different groups: untreated control, 10 mice per inoculum

concentration from 5.10^3 to 5.10^{-1} AFB (n = 50); treated mice, 10 mice for each inoculum ranging from 5.10^3 to 5.10^0 AFB (n = 40) and for each antibiotic: ofloxacin 150 mg/kg; moxifloxacin 150 mg/kg; garenoxacin 150 mg/kg; and clarithromycin 150 mg/kg included as a positive control. Single dose was given the day after inoculation and randomization, and all drugs were administrated by gavage in 0.2 mL sterile water as a single dose of antibiotic.

Assessment of the effectiveness of treatment

Mice were held for 12 months, a period of time sufficient to permit multiplication of a single surviving organism to multiply to a readily countable level. At the end of this period, tissues from the footpad were removed aseptically and homogenized in a final 2 ml volume of Hank's solution as described by Shepard and McRae method [20–22]. *M. leprae* was considered to have multiplied (i.e., viable organisms survived the treatment) in those footpads found to contain $\geq 10^5$ bacilli.

Molecular detection of second-step mutants

Total DNA was extracted from footpad of all mice inoculated with 5.10^3 AFB, following the heat-shock procedure [23]. DNA was subjected to PCR amplifying the QRDRs in *gyrA* as previously described [10] and in *gyrB* using the following primers: GyrBlepS: 5' ACG AGA GTT AGT GCG TCG AAA 3' and GyrBlepAS: 5' GCT GCG CTA AAA ACA CGT AC 3'. Typical reaction mixtures (50 μ l) contained $0.5 \times$ reaction buffer, 2.5 mM of MgCl₂, 0.25 mM of dNTPs, 0.4 μ M of each primer (Eurofins MWG operon), 0.01 U of Taq polymerase (BIO X ACT SHORT TAQ POL, BIOLINE, France) and 5 ml of DNA extract. PCR-amplified fragments were purified by using QIAGEN DNA purification kit (QIAGEN, France) and sequenced by the dideoxy-chain termination method with the ABI PRISM BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Courtaboeuf, France). The oligonucleotide primers used for DNA sequencing were those used for PCR. The nucleotide and deduced amino acid sequences were analyzed with the Seqscape v2.0 software (Applied Biosystems).

Statistical analysis

The proportion of viable *M. leprae* organisms remaining after treatment was determined as the 50% infectious dose, and the significance of the differences between the groups was calculated by the method of Spearman and Kärber [22]. For multiple comparisons between the groups, Bonferroni's correction was applied, i.e., the difference would be significant at the 0.05 level only if the *P* value adjusted to the number of groups: $0.05/n$ in which *n* was defined as the number of primary comparisons. Thus the corrected *P* was $0.05/5 = 0.01$.

Results

Mice survival

Fifty five mice died during the study. Twenty seven mice died due to their advanced age. Twenty eight died due to an accidental problem of water supply during the experiment: 4 mice in ofloxacin 5.10^3 group, 7 mice in untreated control group 5.10^2 , 1 mouse in garenoxacin 5.10^1 group, 1 mouse in garenoxacin 5.10^0 group, 3 mice in clarithromycin 5.10^0 group and 1 mouse in untreated control 5.10^{-1} group.

Bactericidal activity

Results are presented in table 1. In the untreated control group there were 22.6% viable bacilli at the end of the 12 months.

Table 1. Bactericidal activities against *M. leprae* Hoshizuka-4 of ofloxacin, moxifloxacin and garenoxacin measured in mice by the proportional bactericidal method.

Treatment ^a	No. of footpads showing multiplication ^b of <i>M. leprae</i> /no. of footpads harvested, by inoculum ^f					% Viable <i>M. leprae</i> ^c p value ^d	% Viable <i>M. leprae</i> killed by treatment ^e
	5×10^3	5×10^2	5×10^1	5×10^0	5×10^{-1}		
None (control)	6/6	3/3	9/9	5/7	0/9	22.60	
Clarithromycin	7/7	9/9	7/10	0/8		2.17	0.0005
Ofloxacin	1/1	8/8	9/9	1/7		6.06	0.014
Moxifloxacin	6/6	8/8	6/9	1/9		2.16	0.0005
Garenoxacin	6/6	10/10	6/9	1/8		2.7	0.0009

^aAll drugs were administered by gavage as a single dose.

^b*M. leprae* bacilli were considered to have multiplied if the harvest from a footpad yielded $\geq 10^5$ acid-fast bacilli. When the maximum inoculum was 5×10^3 bacilli per footpad, a proportion of viable *M. leprae* as small as 0.006% could be measured.

^cDerived from the following equation: % viable *M. leprae* = $0.69/50 \times$ infectious dose [22].

^dp value (comparison with control group).

^eCalculated from the comparison of the proportions of viable organisms between untreated controls and the treated group.

^fthe high number of dead mice is due to an accidental problem of water supply that occurred during the experiment.

doi:10.1371/journal.pntd.0002559.t001

Clarithromycin killed 90% of viable bacilli, ofloxacin 73%, moxifloxacin 90% and garenoxacin 88%. Compared to untreated control group, the percentage of viable bacilli was significantly smaller in the following treated groups: $p=0.0005$ for clarithromycin, $p=0.0005$ for moxifloxacin, $p=0.0009$ for garenoxacin. For ofloxacin the percentage of viable bacilli was smaller than that of control group but not after Bonferroni correction ($p=0.014$). On the other hand, the percentage of viable bacilli was similar in the group treated by ofloxacin compared to groups treated by moxifloxacin and garenoxacin ($p=0.276$ and 0.334).

Clarithromycin was as active as garenoxacin ($p=0.723$) and moxifloxacin ($p=0.757$). Clarithromycin was more active than ofloxacin but not after Bonferroni correction ($p=0.034$).

Detection of second-step mutants

No mutation in *gyrA* or *gyrB* was found in mice footpads demonstrating the absence of second-step mutant selection in our experiment. This result is not surprising since a single pulse drug is unlikely to result in selection of second step mutations.

Discussion

Phenotypic assessment of *M. leprae* drug resistance is usually done using the continuous method in the mouse footpad model [10,18,24]. This method does not allow assessing bactericidal activity. This study is the first, to the best of our knowledge, assessing *in vivo* the bactericidal activity of various antibiotics of the same family against a *M. leprae* strain carrying mutation involved in drug resistance. Although multiple doses of treatment have been used by others, we chose to treat mice with a single dose of fluoroquinolones in order to be able to compare our present results to our previously published results [25,26]. We demonstrated that despite the presence of a GyrA substitution well known to confer FQ resistance, i.e. A91V [10,13,15], the 3 fluoroquinolones tested had still some *in vivo* activity (Table 1). Ofloxacin activity was marginally significant, but garenoxacin and moxifloxacin remained active. We don't believe that mortality due to the water supply problem biased significantly the ofloxacin results because, for all groups of mice, including the untreated control group, there was no difference between 5.10^2 and 5.10^3 inocula, and all of mice inoculated with 5.10^2 and 5.10^1 and receiving ofloxacin were positive. Therefore it's highly probable that all mice would have been positive in 5.10^3 ofloxacin. Also the rank of activity between ofloxacin and moxifloxacin seen in wild-type strains in a previous study was maintained [27].

Despite the general rule of cross resistance between quinolones, garenoxacin, a new non-fluorinated quinolone, retains most of its activity against strains harboring QRDR mutations, in species such as *Streptococcus pneumoniae* and *Helicobacter pylori* [28–31]. This characteristic, combined with a lower rate of resistant mutant and favorable PK-PD parameters contribute to its higher activity against strains harboring DNA gyrase mutations compared to other quinolones. We previously demonstrated that garenoxacin has the same inhibitory activity against purified *M. leprae* DNA gyrase carrying mutation implicated in quinolone resistance than against the wild-type enzyme [15]. Although less active than moxifloxacin against wild-type strains [32], garenoxacin is as active as moxifloxacin against the GyrA A91V mutant. Garenoxacin is currently under development in several countries [33].

Surprisingly, moxifloxacin also retained most of its activity against the strain harboring the GyrA A91V substitution, despite this substitution reduces moxifloxacin inhibition activity against purified gyrase [15]. Two parameters could explain the maintained activity of moxifloxacin against the strain harbouring the

GyrA A91V substitution. First, it should be kept in mind that moxifloxacin is more active than ofloxacin [25] and garenoxacin [32] against wild-type *M. leprae* [27]. More importantly, it is likely that moxifloxacin MIC against the mutant strain remains lower than moxifloxacin peak serum level, thus allowing some *in vivo* activity as already shown for *M. tuberculosis* [34].

Ofloxacin is naturally less active than moxifloxacin and garenoxacin against *M. leprae* [25,27,32]. In other bacterial species like *M. tuberculosis*, ofloxacin is also less active than moxifloxacin against susceptible strains and this difference remains also against strains harboring DNA gyrase mutations [35]. In other words, in case of DNA gyrase mutation, susceptibility decreases for both ofloxacin and moxifloxacin but the difference of activity between these 2 antibiotics remains the same and explains why despite maintained moxifloxacin activity, ofloxacin was only marginally active against *M. leprae*.

How these activities observed in a murine model can be translated in humans? We used Nude mice (NMRI-*Foxn1*tm/*Foxn1*tm) rather than Swiss or conventional BALB/c because this species is more sensitive for the detection of antibiotic activity [36]. As the strain used was multidrug resistant we could not use dapson nor rifampin as controls and consequently chose clarithromycin. In a previous study conducted in Swiss mice inoculated with a wild-type *M. leprae* strain, the killing rate was lower for clarithromycin (75%) than for moxifloxacin (92%) [25]. In the present study, the killing rates were equivalent between these two antibiotics (Table 1). Thus GyrA A91V substitution reduced the activity of moxifloxacin *in vivo* that became equivalent to that of clarithromycin. In human, clarithromycin although less active than rifampin, has shown bactericidal activity [37]. Thus in case of multidrug resistance, moxifloxacin could still be used in combination with clarithromycin, in a second-line drug scheme against GyrA A91V mutants. We showed a similar phenomenon in *M. tuberculosis* in which the substitution A90V (equivalent to A91V in *M. leprae*) downgrades the bactericidal moxifloxacin into a bacteriostatic drug in immunocompetent Swiss mice [34].

An important point regarding translation of mouse results to human is the dosing of antibiotics used. The 150 mg/kg dosing used generates, in the mouse, an AUC (Area Under the Curve) equivalent to the 400 mg dosing in human for the 3 fluoroquinolones [34,38–42]. For clarithromycin, the 150 mg/kg dosing generates an AUC equivalent to 500 mg twice-daily human dosing [43,44]. Thus, for all tested drugs, the AUC, which is the main PK parameter predicting efficacy, was equivalent to AUC in human at the usual dosing of the antibiotic. Taken together these data indicate that the GyrA A91V substitution confers low-level fluoroquinolone resistance in *M. leprae* and that moxifloxacin can be used in humans against such mutant strains.

Molecular tools are more and more described and used for the diagnosis of drug resistance in leprosy [10,13,19,45]. We demonstrated in the present study that detection of a mutation is not sufficient to exclude a drug from therapeutic regimen, especially when there are a few or no other alternatives. Regarding fluoroquinolones, the present study is relevant for leprosy since, in *M. leprae*, the GyrA A91V substitution is the most prominent substitution described in the literature [13,19,45,46], while the other substitution found in *M. leprae* GyrA (G89C substitution, corresponding to G88C in the *M. tuberculosis* numbering system) has been described only once [13]. Based on results from enzymatic studies performed on *M. leprae* and *M. tuberculosis*, the latter substitution should decrease susceptibility to fluoroquinolones at least at the level obtained with GyrA A91V substitution and could lead to a high-level resistance phenotype [35]. Thus the drug-resistance level generated by DNA gyrase mutation probably

differs depending on the mutations and the therapeutic consequences should also differ.

Equivalent *in vivo* experiments with *rpoB* and *folP* mutants would be important in order to measure the *in vivo* impact of drug resistance on rifampin and dapsone activity. Regarding rifampin, low-level resistance has been extensively studied in *M. tuberculosis*. For example, *M. tuberculosis* strains harboring *rpoB* L533P mutation display low-level rifampin resistance and the same mutation described in *M. leprae* [13] would deserve further evaluation.

References

- Scollard DM, Adams LB, Gillis TP, Krahenbuhl JL, Truman RW, et al. (2006) The continuing challenges of leprosy. *Clin Microbiol Rev* 19: 338–381.
- World Health Organisation (2012) Global leprosy situation, 2012. WHO 87: 317–328.
- Richardus JH, Habbema JDF (2007) The impact of leprosy control on the transmission of *M. leprae*: is elimination being attained? *Lepr Rev* 78: 330–337.
- You E-Y, Kang TJ, Kim S-K, Lee S-B, Chae G-T (2005) Mutations in genes related to drug resistance in *Mycobacterium leprae* isolates from leprosy patients in Korea. *J Infect* 50: 6–11.
- Matsuoka M, Budiawan T, Aye KS, Kyaw K, Tan EV, et al. (2007) The frequency of drug resistance mutations in *Mycobacterium leprae* isolates in untreated and relapsed leprosy patients from Myanmar, Indonesia and the Philippines. *Lepr Rev* 78: 343–352.
- Guelpa-Lauras CC, Cartel JL, Constant-Desportes M, Millan J, Bobin P, et al. (1987) Primary and secondary dapsone resistance of *M. leprae* in Martinique, Guadeloupe, New Caledonia, Tahiti, Senegal, and Paris between 1980 and 1985. *Int J Lepr Other Mycobact Dis* 55: 672–679.
- Chen JK, Wang SY, Hou YH, Ni GX, Zhang JL, et al. (1989) Primary dapsone resistance in China. *Lepr Rev* 60: 263–266.
- Grosset JH, Guelpa-Lauras CC, Bobin P, Brucker G, Cartel JL, et al. (1989) Study of 39 documented relapses of multibacillary leprosy after treatment with rifampin. *Int J Lepr Other Mycobact Dis* 57: 607–614.
- Ji B, Grosset J (2000) Combination of rifapentine-moxifloxacin-minocycline (PMM) for the treatment of leprosy. *Lepr Rev* 71 Suppl: S81–87.
- Cambau E, Bonnafous P, Perani E, Sougakoff W, Ji B, et al. (2002) Molecular Detection of Rifampin and Ofloxacin Resistance for Patients Who Experience Relapse of Multibacillary Leprosy. *Clin Infect Dis* 34: 39–45.
- Pardillo FEF, Burgos J, Fajardo TT, Dela Cruz E, Abalos RM, et al. (2008) Powerful bactericidal activity of moxifloxacin in human leprosy. *Antimicrob Agents Chemother* 52: 3113–3117.
- Cambau E, Perani E, Guillemin I, Jamet P, Ji B (1997) Multidrug-resistance to dapsone, rifampicin, and ofloxacin in *Mycobacterium leprae*. *Lancet* 349: 103–104.
- Maeda S, Matsuoka M, Nakata N, Kai M, Maeda Y, et al. (2001) Multidrug Resistant *Mycobacterium leprae* from Patients with Leprosy. *Antimicrob Agents Chemother* 45: 3635–3639.
- Kim S-K, Lee S-B, Kang T-J, Chae G-T (2003) Detection of gene mutations related with drug resistance in *Mycobacterium leprae* from leprosy patients using Touch-Down (TD) PCR. *FEMS Immunol Med Microbiol* 36: 27–32.
- Matrat S, Cambau E, Jarlier V, Aubry A (2008) Are All the DNA Gyrase Mutations Found in *Mycobacterium leprae* Clinical Strains Involved in Resistance to Fluoroquinolones? *Antimicrob Agents Chemother* 52: 745–747.
- Yokoyama K, Kim H, Mukai T, Matsuoka M, Nakajima C, et al. (2012) Impact of amino acid substitutions in B subunit of DNA gyrase in *Mycobacterium leprae* on fluoroquinolone resistance. *PLoS Negl Trop Dis* 6: e1838.
- Yokoyama K, Kim H, Mukai T, Matsuoka M, Nakajima C, et al. (2012) Amino acid substitutions at position 95 in GyrA can add fluoroquinolone resistance to *Mycobacterium leprae*. *Antimicrob Agents Chemother* 56: 697–702.
- Matsuoka M, Kashiwabara Y, Liangfen Z, Goto M, Kitajima S (2003) A Second Case of Multidrug-resistant *Mycobacterium leprae* Isolated from a Japanese Patient with Relapsed Lepromatous Leprosy. *International Journal of Leprosy and Other Mycobacterial Diseases* 71: 240.
- Da Silva Rocha A, Cunha M das G, Diniz LM, Salgado C, Aires MAP, et al. (2012) Drug and multidrug resistance among *Mycobacterium leprae* isolates from Brazilian relapsed leprosy patients. *J Clin Microbiol* 50: 1912–1917.
- Colston MJ, Hilson GR, Banerjee DK (1978) The “proportional bactericidal test”: a method for assessing bactericidal activity in drugs against *Mycobacterium leprae* in mice. *Lepr Rev* 49: 7–15.
- Shepard CC, McRae DH (1968) A method for counting acid-fast bacteria. *Int J Lepr Other Mycobact Dis* 36: 78–82.
- Shepard CC (1982) Statistical analysis of results obtained by two methods for testing drug activity against *Mycobacterium leprae*. *Int J Lepr Other Mycobact Dis* 50: 96–101.
- Woods SA, Cole ST (1989) A rapid method for the detection of potentially viable *Mycobacterium leprae* in human biopsies: a novel application of PCR. *FEMS Microbiol Lett* 53: 305–309.
- Matsuoka M, Kashiwabara Y, Namisato M (2000) A *Mycobacterium leprae* isolate resistant to dapsone, rifampin, ofloxacin and sparflaxacin. *Int J Lepr Other Mycobact Dis* 68: 452–455.
- Consigny S, Bentoucha A, Bonnafous P, Grosset J, Ji B (2000) Bactericidal activities of HMR 3647, moxifloxacin, and rifapentine against *Mycobacterium leprae* in mice. *Antimicrob Agents Chemother* 44: 2919–2921.
- Ji B, Chauffour A, Andries K, Jarlier V (2006) Bactericidal activities of R207910 and other newer antimicrobial agents against *Mycobacterium leprae* in mice. *Antimicrob Agents Chemother* 50: 1558–1560.
- Matrat S, Petrella S, Cambau E, Sougakoff W, Jarlier V, et al. (2007) Expression and purification of an active form of the *Mycobacterium leprae* DNA gyrase and its inhibition by quinolones. *Antimicrob Agents Chemother* 51: 1643–1648.
- Azoulay-Dupuis E, Bédos JP, Mohler J, Peytavin G, Isturiz R, et al. (2004) Activities of garenoxacin against quinolone-resistant *Streptococcus pneumoniae* strains *in vitro* and in a mouse pneumonia model. *Antimicrob Agents Chemother* 48: 765–773.
- Yamamoto K, Yanagihara K, Sugahara K, Imamura Y, Seki M, et al. (2009) *In vitro* activity of garenoxacin against *Streptococcus pneumoniae* mutants with characterized resistance mechanisms. *Antimicrob Agents Chemother* 53: 3572–3575.
- Suzuki H, Nishizawa T, Muraoka H, Hibi T (2009) Sitaflaxacin and garenoxacin may overcome the antibiotic resistance of *Helicobacter pylori* with *gyrA* mutation. *Antimicrob Agents Chemother* 53: 1720–1721.
- Fukuda Y, Takahata M, Sugiura Y, Shimura Y, Nomura N (2012) *In vitro* pharmacodynamic evaluation of garenoxacin against quinolone-resistant *Streptococcus pneumoniae*. *Int J Antimicrob Agents* 39: 163–167.
- Gidoh M (2007) [In-vitro and in-vivo activities of moxifloxacin and garenoxacin against *Mycobacterium leprae*]. *Nihon Hansenbyo Gakkai Zasshi* 76: 11–17.
- European Medicines Agency (2007) Questions and answers on the withdrawal of the marketing application for Garenoxain Mesylate. London: European Medicines Agency.
- Poissy J, Aubry A, Fernandez C, Lott M-C, Chauffour A, et al. (2010) Should Moxifloxacin Be Used for the Treatment of Extensively Drug-Resistant Tuberculosis? An Answer from a Murine Model. *Antimicrob Agents Chemother* 54: 4765–4771.
- Matrat S, Veziris N, Mayer C, Jarlier V, Truffot-Pernot C, et al. (2006) Functional Analysis of DNA Gyrase Mutant Enzymes Carrying Mutations at Position 88 in the A Subunit Found in Clinical Strains of *Mycobacterium tuberculosis* Resistant to Fluoroquinolones. *Antimicrob Agents Chemother* 50: 4170–4173.
- Ji B, Perani EG, Petinom C, Grosset JH (1996) Bactericidal activities of combinations of new drugs against *Mycobacterium leprae* in nude mice. *Antimicrob Agents Chemother* 40: 393–399.
- Ji B, Jamet P, Perani EG, Sow S, Lienhardt C, et al. (1996) Bactericidal activity of single dose of clarithromycin plus minocycline, with or without ofloxacin, against *Mycobacterium leprae* in patients. *Antimicrob Agents Chemother* 40: 2137–2141.
- Stambaugh JJ, Berning SE, Bulpitt AE, Hollender ES, Narita M, et al. (2002) Ofloxacin population pharmacokinetics in patients with tuberculosis. *Int J Tuberc Lung Dis* 6: 503–509.
- Grosset JH, Guelpa-Lauras CC, Perani EG, Beoletto C (1988) Activity of ofloxacin against *Mycobacterium leprae* in the mouse. *Int J Lepr Other Mycobact Dis* 56: 259–264.
- Lubasch A, Keller I, Borner K, Koeppe P, Lode H (2000) Comparative pharmacokinetics of ciprofloxacin, gatifloxacin, grepafloxacin, levofloxacin, trovafloxacin, and moxifloxacin after single oral administration in healthy volunteers. *Antimicrob Agents Chemother* 44: 2600–2603.
- Andes D, Craig WA (2003) Pharmacodynamics of the New Des-F(6)-Quinolone Garenoxacin in a Murine Thigh Infection Model. *Antimicrob Agents Chemother* 47: 3935–3941.
- Gajjar DA, Bello A, Ge Z, Christopher L, Grasela DM (2003) Multiple-dose safety and pharmacokinetics of oral garenoxacin in healthy subjects. *Antimicrob Agents Chemother* 47: 2256–2263.
- Davey PG (1991) The pharmacokinetics of clarithromycin and its 14-OH metabolite. *Journal of Hospital Infection* 19: 29–37.
- Tessier PR, Kim M-K, Zhou W, Xuan D, Li C, et al. (2002) Pharmacodynamic Assessment of Clarithromycin in a Murine Model of Pneumococcal Pneumonia. *Antimicrob Agents Chemother* 46: 1425–1434.

Acknowledgments

The authors thank Maureen Beaudoin for technical assistance and Renaud Mozet for computational assistance.

Author Contributions

Conceived and designed the experiments: NV AA VJ. Performed the experiments: AC SH MM. Analyzed the data: NV AA VJ SE. Contributed reagents/materials/analysis tools: AC MM SH. Wrote the paper: NV AA VJ.

45. Cambau E, Chauffour-Nevejans A, Tejmar-Kolar L, Matsuoka M, Jarlier V (2012) Detection of antibiotic resistance in leprosy using GenoType LeptraDR, a novel ready-to-use molecular test. *PLoS Negl Trop Dis* 6: e1739.
46. Matsuoka M, Suzuki Y, Estrada Garcia I, Fafutis-Morris M, Vargas-González A, et al. (2010) Possible Mode of Emergence for Drug-Resistant Leprosy Is Revealed by an Analysis of Samples from Mexico. *Jpn J Infect Dis* 63: 412–416.